MycoKeys

MycoKeys 111: 181–210 (2024)
DOI: 10.3897/mycokeys.111.136202

Research Article

Assigning *Xiuguozhangia* (genus *incertae sedis*) to Pseudoberkleasmiaceae (Dothideomycetes, Pleosporomycetidae, Pleosporales) and introducing *X. broussonetiae* as a novel species

Deecksha Gomdola[1,2,3], Rajesh Jeewon[4,5], Eleni Gentekaki[6], Ruvishika S. Jayawardena[2,3], Kevin D. Hyde[3,7], Fatimah Alotibi[7], Yong Wang[1]

1 *Department of Plant Pathology, College of Agriculture, Guizhou University, Guiyang, Guizhou 550025, China*
2 *School of Science, Mae Fah Luang University, Chiang Rai 57100, Thailand*
3 *Center of Excellence in Fungal Research, Mae Fah Luang University, Chiang Rai 57100, Thailand*
4 *Department of Health Sciences, Faculty of Medicine and Health Sciences, University of Mauritius, Réduit 80837, Mauritius*
5 *Department of Zoology, College of Science, King Saud University, P.O. Box 22452, Riyadh 11495, Saudi Arabia*
6 *Department of Veterinary Medicine, University of Nicosia School of Veterinary Medicine, Cyprus 24005, Cyprus*
7 *Department of Botany and Microbiology, College of Science, King Saud University, P.O. Box 22452, Riyadh 11495, Saudi Arabia*
Corresponding author: Yong Wang (yongwangbis@aliyun.com)

OPEN ACCESS

This article is part of:
Exploring the Hidden Fungal Diversity: Biodiversity, Taxonomy, and Phylogeny of Saprobic Fungi
Edited by Samantha C. Karunarathna, Danushka Sandaruwan Tennakoon, Ajay Kumar Gautam

Academic editor:
Samantha C. Karunarathna
Received: 3 September 2024
Accepted: 25 November 2024
Published: 20 December 2024

Citation: Gomdola D, Jeewon R, Gentekaki E, S. Jayawardena R, Hyde KD, Alotibi F, Wang Y (2024) Assigning *Xiuguozhangia* (genus *incertae sedis*) to Pseudoberkleasmiaceae (Dothideomycetes, Pleosporomycetidae, Pleosporales) and introducing *X. broussonetiae* as a novel species. MycoKeys 111: 181–210. https://doi.org/10.3897/mycokeys.111.136202

Copyright: © Deecksha Gomdola et al. This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Abstract

Xiuguozhangia species are dematiaceous hyphomycetes that are characterised by acropleurogenous, dictyoseptate, campanulate or cheiroid, and brown to dark brown conidia that are composed of several layers of cells radiating from a protuberant basal cell, and mostly seen with appendages arising from the apical cells. The genus was introduced based on morphology to accommodate five of the six *Piricaudiopsis* species that exhibited holoblastic conidial ontogeny. *Xiuguozhangia* was referred to as Ascomycota genus *incertae sedis* as it was challenging to resolve its taxonomic placement based solely on the available morphological data (no DNA sequence data was previously available). In this study, we provide DNA sequence data for LSU, ITS, SSU, *TEF1*, and *RPB2* for our isolates, collected from *Broussonetia papyrifera* (Moraceae) in northern Thailand. Based on morphology, we classify our isolates as *Xiuguozhangia*. Since they form a sister lineage to *Pseudoberkleasmium*, we place *Xiuguozhangia* in Pseudoberkleasmiaceae (Pleosporales). Within *Xiuguozhangia*, we establish these two isolates as a new taxon, *Xiuguozhangia broussonetiae*, in view of the presence of new conidiogenous cells developing from subtending cells. *Xiuguozhangia broussonetiae* differs from the extant species in the genus as it has longer conidiophores that are sometimes branched, comprising numerous septa, and its appendages are mostly untapered (sometimes tapering) towards the tips, a feature not observed in other *Xiuguozhangia* species. This is the first study to provide DNA sequence data and phylogenetic relationships for *Xiuguozhangia*. Furthermore, we analysed selected DNA sequence data and provided an updated phylogenetic tree incorporating all families (with representative genera) of Pleosporales.

Key words: 1 new taxon, hyphomycetes, morphology, phylogeny, sequence data, taxonomy

Introduction

Xiuguozhangia was introduced by Zhang et al. (2014) based on morphological characteristics and typified with *X. rosae*. The genus is characterised by effuse, hairy, and brown to dark brown colonies on natural substrates. Conidiophores are erect, straight or flexuous, multiseptate, and brown to dark brown at the base, becoming pale brown towards the apex. Conidiogenous cells are monoblastic, terminal or integrated, lageniform, and pale brown, displaying zero to several percurrent proliferations. In addition, their conidia are acropleurogenous, dictyoseptate, campanulate, cheiroid and brown, comprising multiple layers of cells, either with or without appendages (Zhang et al. 2014; Liu et al. 2024b).

Initially, five *Xiuguozhangia* species were established, all of which are combinations of *Piricaudiopsis viz*., *Xiuguozhangia appendiculata*, *X. indica*, *X. punicae*, *X. rhaphidophorae* and *X. rosae*. *Piricaudiopsis* was established by Mena and Mercado (1987) based on morphology and typified with *P. elegans*. The new combinations of *Xiuguozhangia* were proposed based on their holoblastic conidial ontogeny; *Piricaudiopsis* was reported to have enteroblastic conidial ontogeny (Zhang et al. 2014; Dubey and Jash 2024). However, this morphological difference based on conidial ontogeny is subtle, necessitating careful consideration before applying this trait in generic delineation. Given the limited taxon sampling and analysis across both genera, coupled with the lack of DNA sequence data, it is still uncertain whether *Xiuguozhangia* is actually distinct from *Piricaudiopsis*, or if they could represent the same genus. Both genera were referred to as Ascomycota genera *incertae sedis* in the latest outlines by Wijayawardene et al. (2022a) and Hyde et al. (2024c) as it was difficult to establish their precise taxonomic placement without the availability of DNA sequence data. Recently, based on morphological description, a new species was added to *Xiuguozhangia viz*., *X. macrospora*, collected from dead bamboo stems in India (Dubey and Jash 2024). To date, all extant species of *Xiuguozhangia* have been described as saprobes and collected from China and India (Zhang et al. 2014; Dubey and Jash 2024), while no studies have reported its occurrence in Thailand.

The highlights of this study are as follows: (i) We provide DNA sequence data for multiple gene regions (LSU, ITS, SSU, *TEF1*, and *RPB2*) for our isolates, collected from *Broussonetia papyrifera* (Moraceae) in northern Thailand; (ii) Based on morphology, we classify our isolates as *Xiuguozhangia*. Since they form a sister lineage to *Pseudoberkleasmium*, we place *Xiuguozhangia* in Pseudoberkleasmiaceae (Pleosporales); (iii) Within *Xiuguozhangia*, these two isolates could potentially represent a new taxon, *Xiuguozhangia broussonetiae* in view of the presence of new conidiogenous cells developing from subtending cells. This is the first study to provide DNA sequence data and phylogenetic relationships for *Xiuguozhangia*. In addition, we analysed selected sequence data from GenBank and provided an updated phylogenetic tree incorporating all families (with representative genera) of Pleosporales.

Materials and methods

Sample collection and examination, material deposition, and species delimitation methods

Decaying stems of *Broussonetia papyrifera* (Moraceae) colonised by hyphomycetes were collected from deciduous forest in the Mae Fah Luang University Botanical Garden, Chiang Rai, Thailand during the dry, hot season; collection information was noted (Rathnayaka et al. 2024) and brought to the laboratory in paper boxes for further characterisation (Senanayake et al. 2020). Morphological characteristics were observed using a Motic SMZ 168 Series stereo-microscope. Several conidia were picked from the colonies on the substrate using a sterile needle and placed directly on fresh potato dextrose agar plates (PDA, 39 g/L) containing antibiotics (Amoxicillin, MacroPhar). Conidium germination was observed within 48 hours, and pure cultures were incubated for four weeks at 25 °C under dark conditions. Digital images of micro-morphological characters were captured using a Cannon 750D camera (Canon, Tokyo, Japan) attached to a Nikon ECLIPSE E600 compound microscope (Nikon, Tokyo, Japan) based on the bright-field microscopy technique. Photo plates were assembled using Adobe Photoshop CS6 version 2020 (Adobe Systems, USA), and measurements were made using Tarosoft® Image Frame Work (version 0.97).

The holotype specimen and ex-type living culture have been deposited in the Mae Fah Luang University Herbarium (MFLU) and Mae Fah Luang University Culture Collection (MFLUCC), respectively. Faces of Fungi and Index Fungorum numbers are given for the new species (Jayasiri et al. 2015; Index Fungorum 2024). The description and illustration of the new species have also been updated in the GMS microfungi database (https://gmsmicrofungi.org/) (Chaiwan et al. 2021).

The new species is established based on the morphological species concept and complemented with the phylogenetic species concept (Maharachchikumbura et al. 2021; Pem et al. 2021). Features of *Xiuguozhangia* species are compiled, and a comparison is done to showcase the major differences across existing taxa.

DNA extraction, PCR amplification, and sequencing

Total genomic DNA was extracted from axenic cultures grown on PDA for 28 days using the BIOMIGA Fungus Genomic DNA Extraction Kit (BIOMIGA, San Diego, CA, USA), following the manufacturer's instructions. The internal transcribed spacer (ITS), large subunit (LSU), and small subunit (SSU), and the genes for RNA polymerase 2 (*RPB2*) and translation elongation factor 1α (*TEF1*) were amplified using the following primers: ITS1/ITS4, LR0R/LR5, NS1/NS4 for ITS, LSU, and SSU, respectively (White et al. 1990); fRpb2-5F/7CR for *RPB2* (Liu et al. 1999); and 728F/2218R for *TEF1* (O'Donnell et al. 1998; Carbone and Kohn 1999).

The polymerase chain reaction (PCR) mixture had a final volume of 20 µL, comprising 10 µL of PCR master mix, 1 µL of the forward and reverse primers each (10 µM stock concentration), 7 µL of double-distilled water, and 1 µL of the template DNA. The PCR conditions were as follows: initial denaturation at 95 °C for 3 min; denaturation at 95 °C for 45 s; annealing at 55 °C for 50 s (ITS),

52 °C for 50 s (LSU and SSU), 58 °C for 1 min 30 s (*RPB2* and *TEF1*); extension at 72 °C for 2 min; and final extension at 72 °C for 10 min (number of cycles = 40). Purification and bidirectional sequencing of PCR amplicons were carried out at Sangon Biotech (Shanghai) Co., Ltd., China.

Phylogenetic analyses

The raw reads were checked using DNA Baser Assembler, and ambiguous bases from the 5' and 3' ends were trimmed manually. Consensus sequences were generated using SeqMan (DNAStar, Madison, Wisconsin, USA). The sequences have been deposited in the NCBI GenBank database, and accession numbers for all strains are provided (Table 1). Newly obtained sequences were subjected to blast searches in NCBI, and sequences of ITS, LSU, SSU, *RPB2*, and *TEF1* from other species were retrieved from GenBank (Table 1). Two different datasets were analysed in this study. The first dataset (Dataset 1) evaluated familial relationships and was based on a larger taxon sampling, incorporating representative genera with DNA sequence data from 92 families of Pleosporales. Diademaceae and Lizoniaceae, which also belong to Pleosporales, lack molecular data. Another dataset (Dataset 2) evaluated the phylogenetic relationships within and between *Xiuguozhangia* (Pseudoberkleasmiaceae) and its phylogenetically closely related genera, *Pseudoberkleasmium* (Pseudoberkleasmiaceae) and *Hermatomyces* (Hermatomycetaceae). Dataset 2 was based on multiple strains for these genera.

Single gene datasets were aligned using MAFFT version 7 by applying the default setting (https://mafft.cbrc.jp/alignment/server/) (Katoh et al. 2019) and trimmed using trimAl (Capella-Gutiérrez et al. 2009). The trimmed datasets were concatenated using SequenceMatrix (Vaidya et al. 2011). Maximum likelihood (ML) phylogeny was conducted in the IQ-TREE webserver (https://iqtree.cibiv.univie.ac.at) using the default parameters and 1000 ultrafast bootstrap replicates (Nguyen et al. 2015). The nucleotide substitution model for each DNA marker was automatically generated. The Bayesian information criterion (BIC) selection results were as follows: Dataset 1 – invgamma for ITS, LSU, *RPB2* and *TEF1*, and gamma for SSU; Dataset 2 – gamma for ITS and SSU, and invgamma for LSU, *RPB2* and *TEF1*.

Bayesian inference (BI) was carried out in MrBayes on XSEDE (version 3.2.7a) in the online CIPRES Portal (https://www.phylo.org/portal2) (Huelsenbeck et al. 2001; Ronquist and Huelsenbeck 2003; Miller et al. 2010). Markov chain Monte Carlo (MCMC) sampling was applied to obtain posterior probabilities (PP). Four Markov chains were run simultaneously for 50,000,000 and 5,000,000 generations for datasets 1 and 2, respectively, with trees sampled every 100th generation. Burn-in was set to 20% and the remaining 80% were used to compute the PP of the consensus trees. Phylogenetic trees were visualised in FigTree version 1.4.4 (Rambaut and Drummond 2014).

Genetic distances

To corroborate the phylogenetic placement and evolutionary relationships of the new taxon, intra- and inter-generic genetic distances were computed in MEGA-X by applying the Kimura 2-parameter substitution model, gamma distribution, and pairwise deletion options (Tamura et al. 2013).

Table 1. Names, isolate numbers, and the corresponding GenBank accession numbers of taxa used in the phylogenetic analyses. Type, ex-type, and reference strains are denoted with ᵀ. The new isolates are in bold font.

Species	Isolate/strain number	LSU	ITS	SSU	*TEF1*	*RPB2*
Acrocalymma aquatica	MFLUCC 11-0208 ᵀ	JX276952	JX276951	JX276953	–	–
Acrocalymma pterocarpi	MFLUCC 17 0926 ᵀ	MK347949	MK347732	MK347840	MK360040	–
Ageratinicola kunmingensis	KUMCC 21-0217 ᵀ	NG_243113	NR_191196	NG_242816	–	–
Aigialus grandis	BCC 20000 ᵀ	GU479775	–	GU479739	GU479839	–
Alternaria atrobrunnea	FMR 16868 ᵀ	–	LR537033	–	LR537051	LR537044
Alternaria alternata	AFTOL-ID 1610 ᵀ	DQ678082	KF465761	KC584507	KC584634	KC584375
Amniculicola aquatica	MFLUCC 16-1123 ᵀ	MK106096	–	MK106108	MK109800	–
Amorocoelophoma cassiae	MFLUCC 17-2283	MK347956	MK347739	NG_065775	MK360041	MK434894
Anastomitrabeculia didymospora	MFLU 20-0694 ᵀ	MW412978	NR_172008	NG_073568	MW411338	–
Anastomitrabeculia didymospora	MFLU 11-0236	ON077069	ON077080	ON077074	ON075063	ON075067
Angustimassarina lonicerae	MFLUCC 15-0087	KY496724	KY496759	–	–	–
Anteaglonium gordoniae	MFLUCC 17-2431 ᵀ	MK347977	MK347761	MK347867	MK360042	MK434881
Anteaglonium gordoniae	CD7	–	OK335788	–	–	–
Anteaglonium latirostrum	GKM1119 ᵀ	GQ221874	–	–	GQ221937	–
Anteaglonium latirostrum	GKML100Nb	GQ221876	–	–	GQ221938	–
Aquadictyospora clematidis	MFLUCC 17-2080 ᵀ	MT214545	MT310592	MT226664	MT394727	MT394679
Aquastroma magniostiolata	HHUF 30122 ᵀ	AB807510	LC014540	AB797220	AB808486	–
Aquasubmersa japonica	HHUF 30469 ᵀ	NG_057138	NR_154739	NG_062426	LC194384	LC194421
Aquasubmersa japonica	MFLUCC 17-2121	OP377971	OP377885	OP378047	OP473059	OP473118
Aquasubmersa japonica	MFLUCC 15-0622	OP377958	OP377872	OP378036	OP473051	OP473112
Aquasubmersa mircensis	MFLUCC 11-0401 ᵀ	NG_042699	JX276954	NG_061141	–	–
Aquihelicascus songkhlaensis	MFLUCC 18-1154 ᵀ	MN913692	MT627680	–	MT954380	–
Aquihelicascus songkhlaensis	MFLUCC 18-1273	MN913724	MT627696	MT864319	MT954369	MT878464
Aquimassariosphaeria kunmingensis	KUMCC 18-1019 ᵀ	MT627661	–	MT864312	MT954409	MT878454
Ascocylindrica marina	MD6011 ᵀ	KT252905	–	KT252907	–	–
Ascocylindrica marina	MF416	MK007123	–	MK007124	–	–
Astragalicola vasilyevae	MFLUCC 17-0832 ᵀ	MG828986	NR_157504	MG829098	MG829193	MG829248
Astrosphaeriella fusispora	MFLUCC 10-0555	KT955462	–	KT955443	KT955425	KT955413
Atrocalyx glutinosus	DAOM: 252609 ᵀ	OQ400928	OQ400918	–	OQ413076	OQ413081
Atrocalyx glutinosus	CHEM 2721	OQ400930	OQ400920	–	–	OQ413084
Bahusandhika indica	GUFCC 18001	KF460274	KF460273	–	–	–
Bambusicola bambusae	MFLUCC 11-0614 ᵀ	JX442035	JX442031	JX442039	–	KP761718
Berkleasmium aquaticum (Tubeufiales)	MFLUCC 17-0049 ᵀ	KY790432	KY790444	–	KY792608	MF535268
Berkleasmium aquaticum (Tubeufiales)	MFLUCC 17-0039	KY790431	KY790443	–	KY792607	MF535267
Berkleasmium longisporum (Tubeufiales)	MFLUCC 17-1999 ᵀ	MH558825	MH558698	–	MH550889	MH551012
Berkleasmium longisporum (Tubeufiales)	MFLUCC 17-2002	MH558826	MH558699	–	MH550890	MH551013
Bertiella fici	MFLU 19-2713 ᵀ	MW063223	–	MW079351	MW183786	–
Bertiella fici	NCYU 19-0073	MW063224	–	MW079352	MW183787	–
Bipolaris adikaramae	HSF070 ᵀ	–	MN535176	–	MT548605	–
Boeremia linicola	CBS 116.76 ᵀ	GU237938	GU237754	–	KY484705	KT389574
Boeremia linicola	CBS 248.38	KT389703	KT389486	–	–	KT389575
Biatriospora borsei	NFCCI-4245 ᵀ	MK358813	MK358818	MK358811	MK330938	–
Biatriospora marina	CY 1228	GQ925848	–	GQ925835	GU479848	GU479823
Brevicollum hyalosporum	MAFF 243400 ᵀ	LC271239	LC271242	LC271236	LC271245	LC271249
Brevicollum hyalosporum	MFLUCC 17-0071	MG602200	MG602204	MG602202	MG739516	–
Brevicollum versicolor	HHUF 30591 ᵀ	NG_058716	NR_156335	NG_065124	LC271246	LC271250
Brunneoclavispora camporesii	MFLUCC 11-0001 ᵀ	MN809328	MN809329	–	–	–
Brunneofusispora clematidis	MFLUCC 17-2070 ᵀ	MT214570	MT310615	MT226685	MT394629	MT394692
Camarosporium quaternatum	CPC 31081 ᵀ	NG_064442	NR_159756	KY929123	KY929201	–
Camarosporomyces flavigenus	CBS 314.80 ᵀ	GU238076	MH861266	NG_061093	–	–
Camarosporidiella caraganicola	MFLUCCC 14-0605 ᵀ	KP711381	KP711380	KP711382	–	–
Camarosporidiella melnikii	MFLUCC 17-0684 ᵀ	MF434250	MF434162	MF434338	MF434425	–
Capulatispora sagittiformis	HHUF 29754 ᵀ	NG_042319	NR_119393	NG_060997	LC001756	–
Caryospora aquatica	MFLU 11-1083 ᵀ	NG_059058	NR_156408	MH057850	–	–
Caryospora submersa	MFLUCC 18-1283 ᵀ	MN913720	–	–	–	–
Clematidis italica	MFLUCC 15-0084 ᵀ	KU842381	KU842380	KU842382	–	–
Coelodictyosporium rosarum	MFLUCC 17-0776 ᵀ	MG828991	MG828875	MG829102	MG829195	–
Corylicola italica	MFLU 19-0500 ᵀ	MT554926	MT554925	MT554923	–	MT590776

Species	Isolate/strain number	LSU	ITS	SSU	*TEF1*	*RPB2*
Corynespora cassiicola	CBS 100822	GU301808	–	GU296144	GU349052	GU371742
Corynespora torulosa	CPC 15989 ^{T}	KF777207	NR_145181	–	–	–
Crassimassarina macrospora	MAFF 239606 ^{T}	LC194344	LC194478	LC194302	LC194389	LC194426
Crassimassarina macrospora	HHUF 30512	LC194343	LC194477	LC194301	LC194388	LC194425
Crassiperidium octosporum	MAFF 242971 ^{T}	LC373108	LC373096	LC373084	LC373120	LC373132
Crassiperidium octosporum	MAFF 246401	LC373111	LC373099	LC373087	LC373123	LC373135
Cryptocoryneum japonicum	HHUF 30482 ^{T}	NG_059035	NR_153938	NG_065118	LC096144	LC194438
Cryptocoryneum pseudorilstonei	CBS 113641 ^{T}	NG_059036	NR_153941	LC194322	LC096152	LC194446
Cucurbitaria berberidis	MFLUCC 11-0387	KC506796	–	KC506800	–	–
Curvularia austriaca	CBS 102694 ^{T}	–	MN688802	–	MN688856	–
Curvularia eleusinicola	USJCC-0005 ^{T}	–	MT262877	–	MT432925	–
Cylindroaseptospora leucaenicola	MFLUCC 17-2424 ^{T}	MK347966	NR_163333	MK347856	–	–
Cyclothyriella rubronotata	CBS 141486 ^{T}	KX650544	NR_147651	NG_061252	KX650519	KX650574
Dacampia hookeri	GZU 73897	KT383792	–	–	–	–
Dacampia hookeri	GZU 81840	KT383795	–	–	–	–
Dacampia hookeri	GZU 75980	KT383794	–	–	–	–
Delitschia nypae	MFLUCC 17-2588 ^{T}	–	–	–	MK360049	MK434878
Dendryphion fluminicola	MFLUCC 17-1689 ^{T}	MG208141	NR_157490	–	MG207992	–
Deniquelata cassiae	CMD012A ^{T}	OR500088	OR500092	OR500090	OR501827	–
Dictyocheirospora bannica	KH 332 ^{T}	AB807513	LC014543	AB797223	AB808489	–
Didymella exigua	CBS 183.55 ^{T}	MH868977	MH857436	GU296147	–	–
Didymella rumicicola	CBS 683.79 ^{T}	MH873007	KT389503	–	–	KT389622
Dothidotthia robiniae	MFLUCC 16-1175 ^{T}	MK751817	MK751727	MK751762	MK908017	MK920237
Epicoccum duchesneae	CGMCC 3.18345 ^{T}	KY742249	KY742095	–	–	MT018115
Epicoccum duchesneae	CBS 218.81	MN973322	MN972935	–	–	MN983572
Falciformispora aquatica	MFLUCC 18-0212 ^{T}	MK063643	MK064216	–	MK099811	–
Falciformispora tompkinsii	CBS 200.79 ^{T}	MH872968	MH861199	KF015639	KF015685	KF015719
Fenestella crataegi	CBS 144857 ^{T}	–	NR_165534	–	MK357555	MK357512
Fissuroma calami	MFLUCC 13-0836 ^{T}	MF588993	–	NG_062430	MF588975	–
Flammeascoma lignicola	MFLUCC 10-0128 ^{T}	KT324583	KT324582	KT324584	KT324585	KT324586
Flavomyces fulophazii	CBS 135761 ^{T}	NG_058131	NR_137960	NG_061191	–	–
Foliophoma fallens	CBS 161.78	GU238074	KY940772	GU238215	–	KC584502
Fuscostagonospora cytisi	MFLUCC 16-0622 ^{T}	KY770978	–	KY770977	KY770979	–
Fuscostagonospora sasae	HHUF 29106 ^{T}	AB807548	AB809636	AB797258	AB808524	–
Fusculina eucalypti	CBS 120083 ^{T}	DQ923531	DQ923531	–	–	–
Fusculina eucalyptorum	CBS 145083 ^{T}	MK047499	NR_161140	–	–	–
Fusiformispora clematidis	MFLUCC 17-2077 ^{T}	MT214542	MT310589	MT226661	MT394725	MT394677
Gordonomyces mucovaginatus	CMW 22212 ^{T}	JN712552	JN712486	–	–	–
Halojulella avicenniae	BCC 20173 ^{T}	GU371822	–	GU371830	GU371815	GU371786
Helminthosporiella stilbacea	MFLUCC 15-0813 ^{T}	MT928157	MT928159	MT928161	MT928151	–
Hermatomyces amphisporus	CBS 146610 ^{T}	LR812664	LR812664	–	–	–
Hermatomyces amphisporus	CBS 146613	LR812662	LR812662	–	LR812657	LR812668
Hermatomyces amphisporus	CBS 146614	LR812666	LR812666	–	LR812660	LR812671
Hermatomyces anomianthi	MFLUCC 21-0202 ^{T}	OK655817	OL413437	–	OM117546	–
Hermatomyces bifurcatus	CCF 5900 ^{T}	LS398263	LS398263	–	LS398417	LS398344
Hermatomyces bifurcatus	CCF 5899	LS398262	LS398262	–	LS398416	LS398343
Hermatomyces clematidis	MFLUCC 17-2085 ^{T}	MT214556	MT310603	MT226673	MT394735	MT394684
Hermatomyces constrictus	CCF 5904 ^{T}	LS398264	LS398264	–	LS398418	LS398345
Hermatomyces hainanensis	GZCC 23-0592 ^{T}	OR091329	OR098708	–	–	–
Hermatomyces indicus (=*H. thailandicus*)	MFLUCC 14-1143 ^{T}	KU764692	KU144920	KU712468	–	KU712488
Hermatomyces indicus (=*H. thailandicus*)	MFLUCC 14-1144	KU764693	KU144921	KU712469	–	KU712489
Hermatomyces indicus (=*H. thailandicus*)	MFLUCC 14-1145	KU764694	KU144922	KU712470	KU872756	KU712490
Hermatomyces iriomotensis	KH 361	LC194367	LC194483	–	LC194394	LC194449
Hermatomyces jinghaensis	HKAS 112167 ^{T}	MW989519	MW989495	–	MZ042642	–
Hermatomyces krabiensis (=*H. chiangmaiensis*)	MFLUCC 16-0249 ^{T}	KX525742	KX525750	–	–	KX525754
Hermatomyces krabiensis (=*H. chiangmaiensis*)	MFLUCC 16-2819	KY559394	–	–	–	–
Hermatomyces maharashtraensis	NFCCI 4879 ^{T}	NG_241939	NR_189384	–	MZ130659	MZ130660
Hermatomyces megasporus	CCF 5898 ^{T}	LS398266	LS398266	–	LS398420	–
Hermatomyces megasporus	CCF 5897	–	LS398265	–	LS398419	LS398346
Hermatomyces nabanheensis	KUMCC 16-0149 ^{T}	KY766059	KY766058	KY766060	KY766061	–
Hermatomyces pyriformis	CGMCC 3.27462 ^{T}	PP491962	PP491964	–	PP505452	PP505454
Hermatomyces pyriformis	UESTCC 23.0441	PP491963	PP491965	–	PP505453	PP505455

Species	Isolate/strain number	LSU	ITS	SSU	*TEF1*	*RPB2*
Hermatomyces reticulatus (=*H. subiculosus*)	MFLUCC 15-0843 [T]	KX259523	KX259521	KX259525	KX259527	KX259529
Hermatomyces reticulatus (=*H. subiculosus*)	CCF 5893	LS398267	LS398267	–	LS398421	LS398347
Hermatomyces sphaericoides	CCF 5908 [T]	LS398273	LS398273	–	LS398427	LS398352
Hermatomyces sphaericoides	CCF 5895	LS398270	LS398270	–	LS398424	LS398350
Hermatomyces sphaericus	PMA 116080	LS398281	LS398281	–	LS398431	LS398356
Hermatomyces sphaericus	PMA 116081	LS398283	LS398283	–	LS398432	LS398357
Hermatomyces sphaericus	PRC 4105	–	LS398286	–	–	–
Hermatomyces sphaericus	PRC 4104	–	LS398278	–	LS398430	LS398355
Hermatomyces sphaericus	KZP 462	–	LS398287	–	LS398434	LS398359
Hermatomyces sphaericus	PRM 946201	–	LS398284	–	LS398433	LS398358
Hermatomyces sphaericus	PRC 4116	–	LS398275	–	–	–
Hermatomyces sphaericus	PRC 4100	LS398277	LS398277	–	LS398429	LS398354
Hermatomyces sphaericus	PRC 4106	LS398279	LS398279	–	–	–
Hermatomyces sphaericus	PMA 116085	–	LS398280	–	–	–
Hermatomyces sphaericus	PMA 116082	–	LS398285	–	–	–
Hermatomyces sphaericus	PRC 4117	–	LS398276	–	–	–
Hermatomyces sphaericus	MFLUCC 17-0373	OL782061	OL782144	OL780526	–	–
Hermatomyces sphaericus	HKAS 112725	MW989516	MW989492	–	MZ042639	MZ042636
Hermatomyces sphaericus	HKAS 112166	MW989517	MW989493	–	MZ042640	MZ042637
Hermatomyces sphaericus (=*H. biconisporus*)	KUMCC 17-0183	MH260296	MH275063	MH260338	MH412771	MH412755
Hermatomyces sphaericus (=*H. chromolaenae*)	MFLUCC 16-2818	KY559393	–	–	–	–
Hermatomyces sphaericus (=*H. pandanicola*)	MFLUCC 16-0251	KX525743	KX525751	KX525747	KX525759	KX525755
Hermatomyces sphaericus (=*H. saikhuensis*)	MFLUCC 16-0266	KX525740	KX525748	–	KX525756	KX525752
Hermatomyces sphaericus (=*H. saikhuensis*)	MFLUCC 16-0267	KX525741	KX525749	–	KX525757	KX525753
Hermatomyces sphaericus (=*H. tectonae*)	MFLUCC 14-1140	KU764695	–	NG_063603	KU872757	KU712486
Hermatomyces sphaericus (=*H. tectonae*)	MFLUCC 14-1141	KU764696	KU144918	KU712466	KU872758	–
Hermatomyces sphaericus (=*H. tectonae*)	MFLUCC 14-1142	KU764697	KU144919	KU712467	–	KU712487
Hermatomyces trangensis	BCC 80741 [T]	KY790600	KY790598	KY790602	KY790606	KY790604
Hermatomyces trangensis	BCC 80742	KY790601	KY790599	KY790603	KY790607	KY790605
Hermatomyces tucumanensis	CCF 5912	LS398288	LS398288	–	LS398435	LS398360
Hermatomyces tucumanensis	CCF 5915	LS398290	LS398290	–	LS398437	LS398362
Hermatomyces tucumanensis	CCF 5913	LS398289	LS398289	–	LS398436	LS398361
Hermatomyces turbinatus	MFLUCC 21 0038 [T]	MW989518	MW989494	–	MZ042641	MZ042638
Hermatomyces verrucosus	CCF 5903 [T]	LS398292	LS398292	–	LS398439	LS398364
Hermatomyces verrucosus	CCF 5892	LS398291	LS398291	–	LS398438	LS398363
Hongkongmyces aquaticus	MFLUCC 18-1150 [T]	MN913694	–	MT864302	MT954379	–
Hypsostroma caimitalense	GKM1165 [T]	GU385180	–	–	–	–
Hypsostroma thailandicum	MFLUCC 21-0057 [T]	MZ435867	MZ435865	–	–	–
Jeremyomyces labinae	CBS 144617 [T]	MK442529	MK442589	–	MK442695	MK442665
Juncaceicola alpina	CBS 456.84 [T]	MH873460	MH861761	KY090699	KF253139	KF252188
Keissleriella camporesiana	MFLUCC 15-0029 [T]	MN401741	MN401745	MN401743	MN397907	–
Latorua caligans	CBS 576.65 [T]	NG_058180	–	–	–	–
Latorua grootfonteinensis	CBS 369.72 [T]	NG_058181	–	–	–	–
Lentimurispora urniformis	MFLUCC 18-0497	MH179144	–	MH179160	MH188055	–
Lentithecium clioninum	HHUF 28199 [T]	NG_059391	NR_154137	NG_064845	AB808515	–
Lentithecium pseudoclioninum	HHUF 29055 [T]	NG_059392	AB809633	NG_064847	AB808521	–
Leptosphaeria chatkalica	YGS22 [T]	MW886099	MW886101	MW886100	MW915583	–
Leptosphaerioides guizhouensis	GZAAS 19-4017 [T]	OP099529	OR225067	OR134435	–	–
Leptosphaerioides guizhouensis	GZAAS 19-4018	OP099530	OR225068	OR134436	–	–
Leucaenicola phraeana	MFLUCC 18-0472 [T]	MK348003	MK347785	NG_065784	MK360060	MK434867
Leucaenicola aseptata	MFLUCC 17-2423 [T]	NG_066309	NR_163332	NG_065776	MK360059	MK434891
Libertasomyces myopori	CPC 27354 [T]	NG_058241	KX228281	–	–	–
Ligninsphaeria jonesii	MFLUCC 15-0641 [T]	KU221037	–	–	–	–
Ligninsphaeria jonesii	GZCC 15-0080	KU221038	–	–	–	–
Lindgomyces cigarospora	G619 [T]	KX655804	KX655794	KX655805	–	–
Lindgomyces ingoldianus	ATCC 200398 [T]	AB521736	NR_119938	NG_016531	–	–
Longicorpus striataspora	MFLUCC 18-0267 [T]	MK035988	MK035965	MK035973	MK034428	MK034436
Longiostiolum tectonae	MFLUCC 12-0562 [T]	KU764700	KU712447	KU712459	–	–
Longipedicellata aptrootii	MFLU 10-0297 [T]	KU238894	KU238893	KU238895	KU238892	KU238891
Longipedicellata megafusiformis	SJ-KR4 [T]	MZ538546	MZ538512	–	MZ567090	–
Lonicericola qujingensis	GMBCC1178 [T]	OM855602	OM855593	OM855616	OM857556	–
Lophiostoma carpini	CBS 147279 [T]	MW750386	NR_173000	–	MW752405	MW752384

Species	Isolate/strain number	LSU	ITS	SSU	*TEF1*	*RPB2*
Lophiostoma clavatum	MFLUCC 18-1316 T	MN274566	–	MN304835	MN328901	–
Lophiotrema eburnoides	MAFF 242970 T	LC001707	LC001709	LC001706	LC194403	LC194458
Magnibotryascoma kunmingense	HKAS 111919 T	MW424785	MW424770	MW424800	MW430106	MW430113
Magnibotryascoma rubriostiolatum	CBS 140734 T	–	KU601590	–	KU601609	KU601599
Magnicamarosporium iriomotense	HHUF 30125 T	AB807509	AB809640	AB797219	AB808485	–
Massaria inquinans	CBS 125591 T	MH875187	MH863726	HQ599442	HQ599340	–
Massarina pandanicola	MFLUCC 17-0596 T	MG646947	MG646958	MG646979	MG646986	–
Massarioramusculicola chiangraiensis	MFLUCC 17-2240 T	MH040228	MH040227	MH040229	–	–
Massariosphaeria clematidis	MFLU 16-0174 T	MT214544	MT310591	MT226663	–	–
Matsushimamyces bohaniensis	CBS 140592 T	KR350633	KP765516	–	–	–
Misturatosphaeria aurantonotata	GKM 1238 T	NG_059927	–	–	GU327761	–
Montagnula acaciae	MFLUCC 18-1636 T	ON117298	ON117280	ON117267	ON158093	–
Montagnula aquatica	MFLU 22-0171 T	OP605986	OP605992	OP600504	–	–
Montagnula aquatica	KUNCC 23-14425	OR583116	OR583097	OR583135	OR588088	OR588107
Morosphaeria muthupetensis	NFCC I4219 T	MF614796	MF614795	MF614797	MF614798	–
Multilocularia bambusae	MFLUCC 11-0180 T	KU693438	KU693446	KU693442	–	–
Multiseptospora thailandica	MFLUCC 11-0183 T	KP744490	KP744447	KP753955	KU705657	–
Murilentithecium lonicerae	MFLUCC 18-0675 T	MK214373	MK214370	MK214376	MK214379	–
Murispora galii	MFLUCC 13-0819 T	KT709175	KT736081	KT709182	KT709189	–
Muritestudina chiangraiensis	MFLUCC 17-2551 T	MG602248	MG602247	MG602249	MG602251	MG602250
Neoastrosphaeriella phoenicis	MFLUCC 18-1477 T	MN712339	MN735995	MN699324	MN744232	MN744233
Neobambusicola magnoliae	HKAS 107122 T	ON870389	ON878076	ON870914	–	–
Neocamarosporium goegapense	CPC 23676 T	KJ869220	KJ869163	–	–	–
Neocamarosporium halophilum	RSM70 T	–	OR242722	–	OR289930	–
Neocamarosporium halophilum	BBB RMS57	OR297952	OR297950	–	OR339881	–
Neohelicascus aquaticus	KUMCC 19-0107 T	MT627662	MT627719	MT864314	MT954384	–
Neolophiostoma pigmentatum	MFLUCC 10-0129 T	KT324588	KT324587	KT324589	KT324590	–
Neomassaria fabacearum	MFLUCC 16-1875 T	KX524145	–	NG_061245	KX524149	–
Neomassaria formosana	NTUCC 17-007 T	MH714756	–	MH714759	MH714762	MH714765
Neomassaria hongheensis	KUMCC 21-0344 T	OL423113	OL477614	OL423115	OL754594	OL754595
Neomassarina chromolaenae	MFLUCC 17-1480 T	MT214466	MT214372	MT214419	MT235785	MT235822
Neomassarina pandanicola	MFLUCC 16-0270 T	MG298946	MG298945	MG298947	–	–
Neomassarina thailandica	MFLU 11-0144 T	NG_059718	NR_154244	–	–	–
Neomassarina thailandica	MFLUCC 17-1432	MT214467	MT214373	MT214420	–	–
Neooccultibambusa thailandensis	MFLUCC 16-0274 T	MH260308	MH275074	MH260348	MH412780	MH412758
Neophaeosphaeria phragmiticola	KUMCC 16-0216 T	MG837009	–	NG_065735	MG838020	–
Neophaeosphaeria livistonae	NCYUCC 19-0393 T	OQ437387	OQ437390	OQ437393	–	–
Neoplatysporoides aloes	CPC 36068 T	MN567619	NR_166316	–	–	–
Neopyrenochaeta annellidica	MFLU 11-1105 T	MT183502	MT185538	–	–	–
Neopyrenochaeta cercidis	MFLUCC 18-2089 T	MK347932	MK347718	MK347823	–	MK434908
Neopyrenochaeta chiangraiensis	MFLUCC 13-0881 T	MT183503	MT185539	MT214975	MT454041	–
Neopyrenochaetopsis hominis	UTHSC: DI16-238 T	LN907381	LT592923	–	–	LT593061
Neoroussoella chiangmaiensis	MFLU 22-0205 T	OQ065735	OQ065738	OQ065736	OQ186448	OQ186450
Neothyrostroma encephalarti	CPC 35999 T	MN567613	MN562105	–	MN556831	–
Neotorula aquatica	MFLUCC 15-0342 T	KU500576	KU500569	KU500583	–	–
Neotorula submersa	KUMCC 15-0280 T	KX789217	KX789214	–	–	–
Nigrograna italica	MFLU 23-0139 T	OR538591	OR538590	–	OR531366	OR531365
Occultibambusa bambusae	MFLUCC 13-0855 T	KU863112	KU940123	–	KU940193	KU940170
Occultibambusa jonesii	GZCC 16-0117 T	KY628322	–	KY628324	KY814756	KY814758
Ochraceocephala foeniculi	CBS 145654 T	MN516774	MN516753	MN516743	MN520149	MN520145
Ohleria modesta	CBS 141480	–	KX650563	KX650513	KX650534	KX650583
Ohleria modesta	WU 36870	–	KX650562	–	KX650533	KX650582
Omania hydei	SQUCC 13750 T	MW077155	MW077146	MW077162	MW075772	MW276077
Paraconiothyrium kelleni	CBS 149290 T	NG_229027	NR_185757	OP348926	OP328919	–
Paradictyoarthrinium aquatica	MFLUCC 16-1116 T	NG_064501	NR_158861	–	–	–
Paradictyoarthrinium diffractum	MFLUCC 13-0466	KP744498	KP744455	KP753960	–	KX437764
Paradictyocheirospora tectonae	AMH 10301 T	MW854647	MW854646	–	MW854832	–
Paralophiostoma hysterioides	PUFNI 17617	MT912850	MN582758	MN582762	–	MT926117
Paraleptosphaeria polylepidis	MA 57843 T	–	NR_119469	–	–	–
Paraleptosphaeria polylepidis	APA-2999	MK795717	MK795714	MK795720	MK831009	–
Paramonodictys globosa	HKAS 129169 T	OR091331	OR139016	–	OR494045	OR494048
Parapyrenochaeta protearum	CBS 131315	JQ044453	JQ044434	–	–	LT717683

Species	Isolate/strain number	LSU	ITS	SSU	*TEF1*	*RPB2*
Periconia delonicis	MFLUCC 17-2584 T	NG_068611	–	NG_065770	–	MK434901
Periconia pseudodigitata	KT 1395 T	AB807564	LC014591	AB797274	–	–
Phaeomycocentrospora xinjangensis	CGMCC 3.20479 T	OK256190	OK256193	–	–	–
Phaeoseptum mali	MFLUCC 17-2108 T	MK625197	MK659580	–	MK647990	MK647991
Phaeoseptum terricola	MFLUCC 10-0102 T	MH105779	MH105778	MH105780	MH105781	MH105782
Phaeosphaeria oryzae	CBS 110110 T	KF251689	KF251186	GQ387530	–	KF252193
Pleomonodictys capensis	CBS 968.97 T	KY853521	MH862684	–	–	–
Pleomonodictys capensis	DLUCC 1323	MZ420757	MZ420742	–	–	MZ442696
Pleomonodictys descalsii	FMR 12716 T	KY853522	KY853461	–	–	–
Plenodomus changchunensis	CCMJ 5011 T	OL897174	OL996123	OL984031	–	–
Plenodomus changchunensis	CCMJ 5012	OL966928	OL996124	OL984032	–	OL944508
Pleopunctum clematidis	MFLUCC 17-2091 T	MT214573	MT310618	–	MT394632	MT394693
Pleopunctum thailandicum	MFLUCC 21-0039 T	MZ198896	MZ198894	–	MZ172461	–
Polyschema sclerotigenum	UTHSC DI14-305 T	KP769976	KP769975	–	–	–
Prosthemium alni	MFLUCC 17-0240 T	KY815013	KY797636	–	KY815019	–
Prosthemium intermedium	HHUF 30063 T	AB553778	AB554108	–	–	–
Pseudoasteromassaria aquatica	MFLUCC 18-1397 T	MN913721	MT627674	MT864322	MT954378	–
Pseudoastrosphaeriella longicolla	MFLUCC 11-0171 T	KT955476	–	–	KT955438	KT955420
Pseudoastrosphaeriella thailandensis	MFLUCC 11-0144 T	KT955478	–	KT955457	KT955440	KT955416
Pseudoberkleasmium acaciae	MFLUCC 17-2590 T	NG_066316	NR_163343	NG_065782	MK360073	–
Pseudoberkleasmium chiangmaiense	MFLUCC 17-1809 T	MK131260	MK131259	–	MK131261	–
Pseudoberkleasmium chiangmaiense	MFLU 21-0290	OM065940	OM066271	OM065948	OM102996	OM102997
Pseudoberkleasmium chiangmaiense	MFLUCC 17-2088	MT214585	MT310630	MT226698	MT394643	MT394699
Pseudoberkleasmium chiangmaiense	DLUCC 1655	MZ420759	MZ420744	MZ420749	MZ442693	–
Pseudoberkleasmium chiangraiense	MFLUCC 21-0154 T	OL584200	OL584189	OL606408	OL912943	OL697401
Pseudoberkleasmium chiangraiense	MFLUCC 21-0161	OL584201	OL584190	OL606409	OL912944	OL697402
Pseudoberkleasmium chiangraiense	MFLUCC 21-0162	OL584205	OL584191	OL606410	OL912945	OL697403
Pseudoberkleasmium pandanicola	KUMCC 17-0178 T	MH260304	MH275071	MH260344	–	–
Pseudochaetosphaeronema chiangraiense	MFLU 21-0083 T	MZ457922	MZ457923	–	MZ476770	–
Pseudochaetosphaeronema chiangraiense	UESTCC 23.0065	OR253260	OR253108	–	OR251160	–
Pseudochaetosphaeronema irregulare	CGMCC 3.22458 T	OQ758163	OQ798972	OQ758194	OQ809057	OQ809023
Pseudochaetosphaeronema irregulare	CGMCC 3.22461	OQ758162	OQ798971	OQ758193	OQ809056	OQ809022
Pseudocoleodictyospora sukhothaiensis	MFLUCC 12-0554 T	KU764710	KU712440	NG_062416	–	KU712493
Pseudocoleodictyospora thailandica	MFLUCC 12-0565 T	KU764701	NR_154337	NG_062417	–	KU712494
Pseudocoleophoma heteropanacicola	ZHKUCC 23-0880 T	OR365486	OR365456	–	OR700204	–
Pseudolophiotrema elymicola	HHUF 28984 T	LC194381	LC194505	LC194339	LC194418	LC194473
Pseudomassarina clematidis	MFLU 16-0493 T	MT214586	MT415397	MT226699	MT394644	MT394700
Pseudopyrenochaeta lycopersici	FMR 15746 T	EU754205	NR_103581	NG_062728	–	LT717680
Pseudopyrenochaeta terretris	FMR 15327 T	LT623216	LT623228	–	–	LT623287
Pseudotetraploa rajmachiensis	NFCCI 4618 T	MN937204	MN937222	–	–	–
Pseudoxylomyces elegans	KT 2887	AB807598	LC014593	AB797308	AB808576	–
Profundisphaeria fusiformispora	GZAAS 20-4010 T	–	–	OR134442	OR140432	OR146942
Profundisphaeria fusiformispora	GZAAS 20-4012	OR209667	–	OR134443	OR140433	–
Pyrenochaetopsis leptospora	CBS 101635 T	GQ387627	JF740262	NG_063097	MF795881	LT623282
Pyrenochaetopsis tabarestanensis	IBRC:M 30051 T	KF803343	NR_155636	NG_065034	–	–
Quadricrura bicornis	HHUF 30023 T	AB524613	AB524797	AB524472	AB524828	–
Quercicola fusiformis	MFLUCC 18-0479 T	MK348009	MK347790	MK347898	MK360085	MK434864
Quercicola guttulospora	MFLUCC 18-0481 T	MK348010	MK347791	MK347899	MK360086	–
Quixadomyces cearensis	HUEFS 238438 T	MG970695	NR_160606	–	–	–
Roussoella bambusarum	GMBC 0316 T	ON479892	ON479891	–	ON505017	ON505012
Roussoella guttulata	MFLUCC 20-0102 T	MT734818	MT734821	–	MW022188	MW022187
Rubroshiraia bambusae	HKAS 102255 T	MK804658	MK804678	MK804704	MK819218	–
Rubroshiraia bambusae	HKAS 102256	MK804659	MK804679	MK804705	MK819219	–
Salsuginea phoenicis	MFLU 19-0015 T	MK405280	–	–	MK404650	–
Salsuginea ramicola	KT 2597.2	GU479801	–	GU479768	GU479862	GU479834
Salsuginea rhizophorae	MFLU 18-0540 T	MN017851	–	MN017917	–	–
Seltsamia ulmi	CBS 143002 T	MF795794	MF795794	MF795794	MF795882	MF795836
Septoriella chlamydospora	MFLUCC 15-0177 T	KU163654	KU163658	KU163655	–	–
Septoriella hibernica	CBS 145371 T	MK540036	MK539966	–	–	MK540097
Setoarthopyrenia chromolaenae	MFLUCC 17-1444 T	MT214438	MT214344	MT214392	MT235768	MT235805
Shiraia bambusicola	GZAAS 2.0708 T	KC460982	GQ845414	–	–	–
Shiraia bambusicola	HKAS 102267	MK804657	MK804677	MK804703	MK819217	MK819237

Species	Isolate/strain number	LSU	ITS	SSU	*TEF1*	*RPB2*
Sporormurispora atraphaxidis	MFLUCC 17-0742 ^{T}	MG829083	MG828971	MG829183	–	–
Sporormurispora paulsenii	MFLUCC 17-1957 ^{T}	MK966143	–	MK963075	–	MN023029
Stagonospora forlicesenensis	MFLUCC 15-0054 ^{T}	KX655547	KX655557	KX655552	KX655562	–
Stagonospora imperaticola	MFLUCC 15-0026	KY706133	KY706143	KY706138	KY706146	KY706149
Stemphylium clematidis	MFLUCC 14-0937 ^{T}	MT214583	MT310628	MT226696	–	–
Stemphylium carpobroti	CPC 38637 ^{T}	MW175395	MW175355	–	–	–
Striatiguttula nypae	MFLUCC 18-0265 ^{T}	MK035992	MK035969	MK035977	MK034432	MK034440
Striatiguttula phoenicis	MFLUCC 18-0266 ^{T}	MK035995	MK035972	MK035980	MK034435	MK034442
Subglobosporium tectonae	MFLUCC 12-0393 ^{T}	KU764703	KU712445	KU712464	–	KU712485
Subglobosporium tectonae	MFLUCC 12-0390	KU764702	KU712446	KU712463	–	KU712495
Sublophiostoma thailandica	MFLUCC 11-0207 ^{T}	KX534212	MW136257	KX534218	KX550077	MW088714
Sublophiostoma thailandica	MFLUCC 11-0185	KX534216	MW136275	KX534222	KX550080	MW088718
Submersispora variabilis	MFLUCC 17-2360 ^{T}	MN913682	MT627683	MT864310	–	–
Submersispora variabilis	N-KR15	MZ538561	MZ538527	MZ538575	MZ567103	MZ567114
Sulcatispora acerina	HHUF 30449 ^{T}	LC014610	LC014597	LC014605	LC014615	–
Sulcatispora berchemiae	HHUF 29097 ^{T}	AB807534	AB809635	AB797244	AB808509	–
Sulcosporium thailandica	MFLUCC 12-0004	KT426563	MG520958	KT426564	–	–
Teichospora trabicola	C134 ^{T}	KU601591	KU601591	–	KU601601	KU601600
Tetraploa aquatica	MFLU 19-0995 ^{T}	MT530452	MT530448	–	–	–
Tetraploa aquatica	MFLU 19-0996	MT530453	MT530449	MT530454	–	–
Thyridaria acaciae	CBS 138873 ^{T}	NG_058127	KP004469	–	–	–
Thyridaria broussonetiae	CBS 141481 ^{T}	KX650568	KX650568	KX650515	KX650539	KX650586
Thyrostroma jaczewskii	MFLUCC 18-0787 ^{T}	MK765857	MK765856	MK765858	–	–
Torula camporesii	KUMCC 19-0112 ^{T}	MN507402	MN507400	MN507401	MN507403	MN507404
Torula pluriseptata	MFLUCC 14-0437 ^{T}	KY197855	MN061338	KY197862	KY197875	KY197869
Trematosphaeria grisea	CBS 332.50 ^{T}	NG_057979	NR_132039	NG_062930	KF015698	KF015720
Trematosphaeria pertusa	CBS 122368 ^{T}	NG_057809	NR_132040	FJ201991	KF015701	FJ795476
Tubeufia abundata (Tubeufiales)	MFLUCC 17-2024 ^{T}	MH558894	MH558769	–	MH550961	MH551095
Tubeufia aquatica (Tubeufiales)	MFLUCC 16-1249 ^{T}	KY320539	KY320522	–	KY320556	MH551142
Tubeufia hainanensis (Tubeufiales)	GZCC 22-2015 ^{T}	OR030835	OR030842	–	OR046679	OR046685
Tubeufia hainanensis (Tubeufiales)	GZCC 23-0589	OR066421	OR066414	–	OR058860	OR058853
Tzeanania taiwanensis	NTUCC 17-005 ^{T}	MH461120	MH461123	MH461126	MH461130	MH461128
Tzeanania taiwanensis	NTUCC 17-006	MH461121	MH461124	MH461127	MH461131	MH461129
Wicklowia aquatica	CBS 125634 ^{T}	MH875044	OM322822	GU266232	–	GU371813
Wicklowia fusiformispora	N-KR1 ^{T}	MZ538567	MZ538533	MZ538576	MZ567108	–
Xenomassariosphaeria clematidis	MFLUCC 14-0923 ^{T}	MT214571	MT310616	–	MT394630	–
Xenomassariosphaeria rosae	MFLUCC 15-0179 ^{T}	MG829092	–	MG829192	–	–
Xenopyrenochaetopsis pratorum	CBS 445.81 ^{T}	GU238136	MH861363	NG_062792	–	KT389671
Xiuguozhangia broussonetiae	**MFLUCC 24-0258 ^{T}**	**PQ137419**	**PQ137417**	**PQ137421**	**PQ488461**	**PQ488459**
Xiuguozhangia broussonetiae	**MFLUCC 24-0259**	**PQ137420**	**PQ137418**	**PQ137422**	–	**PQ488460**
Zopfia rosatii	CBS 427.62 ^{T}	NG_066246	NR_160090	–	–	–

– Data unavailable.

Results

Phylogenetic analyses

Blast searches of LSU, ITS, SSU, *TEF1* and *RPB2* sequences indicated that the two isolates are highly similar to various genera in Pleosporales, including *Atrocalyx*, *Hermatomyces*, *Lophiotrema* and *Pseudoberkleasmium*.

Dataset 1 consisted of 4250 characters (LSU = 1–846, ITS = 847–1355, SSU = 1356–2359, *TEF1* = 2360–3257, and *RPB2* = 3258–4250), which was analysed to depict relationships at a higher taxonomic level for *Xiuguozhangia* (Fig. 1). Outgroup taxa were selected from Tubeufiales. The log-likelihood of the consensus tree (Fig. 1) was -143487.839. The average standard deviation of split frequencies at the end of the total MCMC generations converged to 0.0092.



Figure 1. Maximum likelihood analysis (IQ-tree) based on a combined dataset of LSU, ITS, SSU, *TEF1*, and *RPB2* sequences of all families (with representative genera) of Pleosporales. Bootstrap support values (ML ≥ 80%) and Bayesian posterior probabilities (PP ≥ 0.95) are given above the branches or near the nodes as ML/PP. Hyphens (--) indicate bootstrap support values below 80% for ML and posterior probabilities below 0.95. The tree is rooted with *Tubeufia abundata* (MFLUCC 17-2024), *T. aquatica* (MFLUCC 16-1249), *T. hainanensis* (GZCC 22-2015 and GZCC 23-0589), *Berkleasmium aquaticum* (MFLUCC 17-0049 and MFLUCC 17-0039) and *B. longisporum* (MFLUCC 17-1999 and MFLUCC 17-2002) (Tubeufiales). Type, ex-type, and reference strains are denoted with T. The new isolates are in bold font. The different colour blocks indicate the families to which the taxa belong.



Figure 1. Continued.



Figure 1. Continued.



Figure 1. Continued.

Dataset 2 comprised 4285 characters (LSU = 1–833, ITS = 834–1344, SSU = 1345–2350, *TEF1* = 2351–3272, and *RPB2* = 3273–4285), and was used to infer the inter-generic relationships of *Xiuguozhangia*, which comprised multiple strains for each genus (Fig. 2). Based on the results from dataset 1, four


ML/PP
Hermatomyces sphaericus HKAS 112166
Hermatomyces sphaericus MFLUCC 17-0373
Hermatomyces sphaericus PMA 116082
Hermatomyces sphaericus HKAS 112725
Hermatomyces sphaericus PMA 116085
Hermatomyces sphaericus (=H. chromolaenae) MFLUCC 16-2818
Hermatomyces sphaericus PRC 4105
Hermatomyces sphaericus PRM 946201
Hermatomyces sphaericus KZP 462
Hermatomyces sphaericus PMA 116080
Hermatomyces sphaericus PMA 116081
Hermatomyces sphaericus PRC 4100
Hermatomyces sphaericus (=H. saikhuensis) MFLUCC 16-0267
Hermatomyces sphaericus PRC 4117
Hermatomyces sphaericus PRC 4116
Hermatomyces sphaericus (=H. saikhuensis) MFLUCC 16-0266
Hermatomyces sphaericus PRC 4104
Hermatomyces sphaericus (=H. biconisporus) KUMCC 17-0183
Hermatomyces sphaericus (=H. tectonae) MFLUCC 14-1140
Hermatomyces sphaericus (=H. tectonae) MFLUCC 14-1141
Hermatomyces sphaericus (=H. tectonae) MFLUCC 14-1142
Hermatomyces sphaericus (=H. pandanicola) MFLUCC 16-0251
Hermatomyces sphaericus PRC 4106
Hermatomyces indicus (=H. thailandicus) MFLUCC 14-1144
Hermatomyces indicus (=H. thailandicus) MFLUCC 14-1145
Hermatomyces indicus (=H. thailandicus) MFLUCC 14-1143
Hermatomyces amphisporus CBS 146610 T
Hermatomyces amphisporus CBS 146613
Hermatomyces amphisporus CBS 146614
Hermatomyces anomianthi MFLUCC 21-0202 T
Hermatomyces nabanheensis KUMCC 16-0149 T
Hermatomyces turbinatus MFLUCC 21-0038 T
Hermatomyces bifurcatus CCF 5899
Hermatomyces bifurcatus CCF 5900 T
Hermatomyces megasporus CCF 5898 T
Hermatomyces megasporus CCF 5897
Hermatomyces reticulatus (=H. subiculosus) CCF 5893
Hermatomyces reticulatus (=H. subiculosus) MFLUCC 15-0843 T
Hermatomyces hainanensis GZCC 23-0592 T
Hermatomyces tucumanensis CCF 5912
Hermatomyces tucumanensis CCF 5915
Hermatomyces tucumanensis CCF 5913
Hermatomyces sphaericoides CCF 5895
Hermatomyces sphaericoides CCF 5908 T
Hermatomyces constrictus CCF 5904 T
Hermatomyces verrucosus CCF 5892
Hermatomyces verrucosus CCF 5903 T
Hermatomyces krabiensis (=H. chiangmaiensis) MFLUCC 16-0249 T
Hermatomyces krabiensis (=H. chiangmaiensis) MFLUCC 16-2819
Hermatomyces iriomotensis KH 361
Hermatomyces trangensis BCC 8-741 T
Hermatomyces trangensis BCC 80742
Hermatomyces jinghaensis HKAS 112167 T
Hermatomyces clematidis MFLUCC 17-2085 T
Hermatomyces maharashtraensis NFCCI4879 T
Hermatomyces pyriformis CGMCC 3.27462 T
Hermatomyces pyriformis UESTCC 23.0441
Pseudoberkleasmium chiangmaiense DLUCC 1655
Pseudoberkleasmium chiangmaiense MFLUCC 17-1809 T
Pseudoberkleasmium chiangmaiense MFLU 21-0290
Pseudoberkleasmium chiangmaiense MFLUCC 17-2088
Pseudoberkleasmium chiangraiense MFLUCC 21-0161
Pseudoberkleasmium chiangraiense MFLUCC 21-0162
Pseudoberkleasmium chiangraiense MFLUCC 21-0154 T
Pseudoberkleasmium pandanicola KUMCC 17-0178 T
Pseudoberkleasmium acaciae MFLUCC 17-2590 T
Xiuguozhangia broussonetiae MFLUCC 24-0258 T
Xiuguozhangia broussonetiae MFLUCC 24-0259
Anteaglonium latirostrum GKM1119 T
Anteaglonium latirostrum GKML100Nb
Anteaglonium gordoniae MFLUCC 17-2431 T
Anteaglonium gordoniae CD7
0.03
Hematomycetaceae
Pseudoberkleasmiaceae
Outgroups
Anteagloniaceae

Figure 2. Maximum likelihood analysis (IQ-tree) based on the combined LSU, ITS, SSU, *TEF1* and *RPB2* sequences of *Xiuguozhangia*, *Pseudoberkleasmium* and *Hermatomyces*, generated from dataset 2. Bootstrap support values (ML ≥ 80%) and Bayesian posterior probabilities (PP ≥ 0.95) are given above the branches or near the nodes as ML/PP. Hyphens (--) indicate bootstrap support values below 80% for ML and posterior probabilities below 0.95. The tree is rooted with *Anteaglonium gordoniae* (MFLUCC 17-2431 and CD7) and *A. latirostrum* (GKM1119 and GKML100Nb) (Anteagloniaceae, Pleosporales). Type, ex-type, and reference strains are denoted with T. The new isolates are in bold font. The different colour blocks indicate the families to which the taxa belong.

taxa belonging to Anteagloniaceae (Pleosporales) were selected as outgroups as they are phylogenetically closely related to Pseudoberkleasmiaceae and Hermatomycetaceae. The log-likelihood of the consensus tree (Fig. 2) was -17829.366. The average standard deviation of split frequencies at the end of the total MCMC generations converged to 0.0031.

The two strains of *Xiuguozhangia broussonetiae* (MFLUCC 24-0258 and MFLUCC 24-0259) grouped with 100% ML and 1.00 PP support. This subclade formed a separate sister lineage to *Pseudoberkleasmium* with 100% ML and 1.00 PP support (Figs 1, 2). Single and combined gene trees from datasets 1 and 2 indicate that our species is phylogenetically most closely related to *Pseudoberkleasmium*, and its placement in Pseudoberkleasmiaceae is supported by maximum bootstrap support and posterior probability in both combined data trees.

Genetic distances

Based on the generic relationship depicted in Figs 1, 2, we computed the group mean distances between *Hermatomyces* spp. (group 1), *Pseudoberkleasmium* spp. (group 2) and *Xiuguozhangia* spp. (group 3) across ITS and LSU markers. The difference in the genetic distances across both markers is given in Table 2.

Table 2. Group mean genetic distances (%) in *Hermatomyces* spp., *Pseudoberkleasmium* spp., and *Xiuguozhangia* spp. across ITS (511 bp) and LSU (833 bp).

DNA markers	Groups	Group 1: *Hermatomyces* spp.	Group 2: *Pseudoberkleasmium* spp.
ITS	**Group 1:** *Hermatomyces* spp.	0	–
	Group 2: *Pseudoberkleasmium* spp.	9.052	0
	Group 3: *Xiuguozhangia* spp.	10.62834205	13.25008949
LSU	**Group 1:** *Hermatomyces* spp.	0	–
	Group 2: *Pseudoberkleasmium* spp.	3.1615	0
	Group 3: *Xiuguozhangia* spp.	4.1404	3.8669

Taxonomy

Xiuguozhangia broussonetiae **Gomdola, Jayaward. & K.D. Hyde, sp. nov.**
Index Fungorum: IF901943
Facesoffungi Number: FoF16323
Fig. 3

Holotype. MFLU 24-0227.

Etymology. The specific epithet refers to the host genus, *Broussonetia*, from which the species was isolated.

Description. Saprobic on decaying stems of *Broussonetia papyrifera*. Sexual morph not observed. Asexual morph on substrate. Hyphomycetous. ***Colonies*** on the substrate effuse, hairy, olivaceous to dark brown, appearing velvety due to numerous conidiophores. ***Mycelium*** semi-immersed or immersed, composed of septate, branched, smooth, hyaline or pale brown to brown hyphae. ***Conidiophores*** (430–)550–750(–890) µm long ($\bar{x}$ = 681 µm, n = 10), 15–24(–34) µm wide ($\bar{x}$ = 21.8 µm, n = 10) at the base, 11–14(–16) µm wide ($\bar{x}$ = 12.3 µm, n = 10) in the middle, 7–10 µm(–12) wide ($\bar{x}$ = 7.8 µm, n = 10) at the apex, rarely branched, macronematous, mononematous, differentiated, smooth, thick-



Figure 3. *Xiuguozhangia broussonetiae* (MFLUCC 24-0258) **A** substrate (*Broussonetia papyrifera*) **B, C** colonies on the substrate **D–F** conidiophores and attachment of conidia **G–I** conidiogenous cells with percurrent proliferations **J–O, Q, R** conidia bearing appendages **P** top view of a conidium **S** top view of colonies on PDA after 7 days **T** top and reverse colonies on PDA after 28 days. Scale bars: 100 µm (**B, C**); 50 µm (**D–F**); 10 µm (**G–I, P**); 20 µm (**J–O, Q, R**).

walled, erect, straight or flexuous, brown to dark brown, wider at the base and ocasionally conical at the apex, comprising 12–17 septa. ***Conidiogenous cells*** 5–12 µm long × 4–9 µm wide ($\overline{x}$ = 7.4 × 6.2 µm, n = 10), holoblastic, enteroblastic, integrated, smooth-walled, ovoid to ampulliform, hyaline or brown to dark brown, occurring terminally or intercalary on conidiophores, with up to four successive percurrent proliferations, with new conidiogenous cells developing on subtending cells. ***Conidia*** 25–40 µm long × 30–60 µm wide ($\overline{x}$ = 35 × 47 µm, n = 30), width measured between the two extremities of the apices, solitary, dictyospored and cheirospored, fan-shaped to cheiroid, lenticular in edge view, occasionally apically 2–3-lobed, dark brown to olivaceous brown, dictyoseptate, with up to 15 columns of cells radiating from a protuberant basal cell 2–3 µm wide, septa obscured by dark pigmentation, and with 1–3 apical appendages (rarely without appendages). ***Apical appendages*** (2–)6–16(–20) µm long ($\overline{x}$ = 10.9 µm, n = 10), 4–5 µm wide ($\overline{x}$ = 4.4 µm, n = 10) at the base, 4–5 µm wide ($\overline{x}$ = 4.1 µm, n = 10) at the apex, mostly short and untapered, sometimes long and tapering, arising from the sides of the outermost rows of cells of the conidium, pale brown to brown, becoming hyaline and rounded at the tips, and consisting of 1–5 septa.

Culture characteristics. On PDA, colony circular with lobate to crenated margin, reaching 25 mm diam. after 28 days incubated at 25 °C, greyish white to olivaceous brown from center to edge, fluffy, raised to convex, penetrating the media and displaying a dark brown colour around the media.

Material examined. Thailand • Chiang Rai Province, Mae Fah Luang University Botanical Garden, on decaying stems of *Broussonetia papyrifera* (Moraceae), 19 May 2023, D. Gomdola, F2-A (MFLU 24-0227, ***holotype***), ex-type MFLUCC 24-0258.

Additional material examined. Thailand • Chiang Rai Province, Mae Fah Luang University Botanical Garden, on decaying stems of *Broussonetia papyrifera* (Moraceae), 19 May 2023, D. Gomdola, F2-B (MFLU 24-0228), living culture MFLUCC 24-0259.

GenBank accession numbers. MFLUCC 24-0258; ITS = PQ137417, LSU = PQ137419, SSU = PQ137421, *RPB2* = PQ488459 and *TEF1* = PQ488461; MFLUCC 24-0259; ITS = PQ137418, LSU = PQ137420, SSU = PQ137422 and *RPB2* = PQ488460.

Notes. Our two isolates (MFLUCC 24-0258 and MFLUCC 24-0259) group together with 100% ML and 1.00 PP support, and this subclade is sister to *Pseudoberkleasmium* species with 100% ML and 1.00 PP support (Figs 1, 2).

A close comparison of the morphological characters across the existing *Xiuguozhangia* species is collated and presented in Table 3 to support the establishment of the new species, *X. broussonetiae*.

A summary of the main findings from the morphological assessment is presented below:

1. The conidiophores of our species, *Xiuguozhangia broussonetiae*, are longer than all other *Xiuguozhangia* taxa (up to 890 µm long), comprising numerous septa (up to 17). Furthermore, conidiophores of other *Xiuguozhangia* species are unbranched, while those of *X. broussonetiae* are sometimes branched (Fig. 3E).
2. The conidia of *Xiuguozhangia broussonetiae* (Figs 3J–O, Q, R) consist of up to 15 columns of cells radiating from a protuberant basal cell that is smaller (2–3 µm wide) compared to those of other species.

3. The appendages of *X. broussonetiae* are mostly untapered (Figs 3N, 3P–R) and sometimes taper towards the tips (Fig. 3B), whereas all other *Xiuguozhangia* species have only tapering appendages.
4. *Xiuguozhangia broussonetiae* differs from all extant species but has a close morphological overlap with *X. rosae*. However, the primary feature that demarcates *X. broussonetiae* from *X. rosae* is the number and features of the apical appendages. Up to three appendages were observed in *X. broussonetiae* (sometimes seen without appendage), while *X. rosae* has two appendages. In addition, the appendages of *X. broussonetiae* are 1–5-septate, while those of *X. rosae* display one to two septa.
5. *Xiuguozhangia broussonetiae* has percurrently proliferating, enteroblastic conidiogenous cells that sometimes produce new conidiogenous cells on subtending cells (Fig. 3D–I). This feature has not been observed in other species of the genus.

Based on these morphological differences, we conclude that our taxon is distinct from the existing *Xiuguozhangia* species.

Discussion

The number of hyphomycetes introduced over the past decade has increased substantially, indicating that their diversity is rather high (Hyde et al. 2020, 2023, 2024a, b; Bhunjun et al. 2022; Manawasinghe et al. 2022; Calabon et al. 2023; Liu et al. 2024b; Tian et al. 2024; Zhang et al. 2024). Hyphomycetes have a ubiquitous distribution in aquatic and terrestrial habitats, occurring on different substrates in tropical and subtropical regions (Seifert et al. 2011; Bao et al. 2021; Wu and Diao 2022, 2023; Dong et al. 2023; Senanayake et al. 2023; Yang et al. 2023; Liu et al. 2024a, b). Several hyphomycetes are rather speciose, for example, *Cladosporium*, *Dictyosporium*, *Helminthosporium*, and *Sporidesmium* (Shenoy et al. 2006; Hu et al. 2023; Yang et al. 2023). However, the hyphomycete genus, *Xiuguozhangia*, does not appear to be species-rich, and no studies have reported its occurrence in Thailand. This study introduces a new species, *Xiuguozhangia broussonetiae*, from *Broussonetia papyrifera* in northern Thailand.

There are more than 30,000 fungal species that exhibit the asexual morph, belonging to 2265 hyphomycetous genera (Wijayawardene et al. 2022b; Jayawardena et al. 2023; Perera et al. 2023; Ma et al. 2024; Zhang et al. 2024). The *Xiuguozhangia* species reported so far occur only in their asexual morph as saprobic hyphomycetes. A probable explanation for their asexual morph occurrence might be for survival, as conidia can be produced in large amounts and are easily released for dispersal compared to ascospores (Gilbert and Parker 2023). Their sexual morphs have not been reported yet, and no research has been conducted to link their asexual and sexual morphs, possibly due to the lack of molecular data and their availability in cultures. To date, all *Xiuguozhangia* taxa have been documented from Asia (China, India and Thailand – this study). We anticipate discovering more species in this genus, as well as the sexual morph, especially in high biodiversity areas like China and Thailand, as these taxa have not yet been extensively collected, isolated, and studied.

In the phylogenetic analysis of dataset 1, which included representative taxa from all families of Pleosporales, *Xiuguozhangia* forms a sister lineage

Table 3. Phenotypic comparison of *Xiuguozhangia* species. Features that depict *X. broussonetiae* from other *Xiuguozhangia* species are in bold.

Species and characters		*X. appendiculata*	*X. broussonetiae*	*X. indica*	*X. macrospora*	*X. punicae*	*X. rhaphidophorae*	*X. rosae*
Colonies on the natural substrates	**Features**	Effuse, hairy	Effuse, appearing velvety due to numerous conidiophores	Effuse, hairy	Effuse, hairy, growing in association with species of lichen	Effuse, hairy	Effuse, hairy	Effuse, hairy
	Colour	Olivaceous to dark brown	Olivaceous to dark brown	Olivaceous to dark brown	Olivaceous to dark brown	Olivaceous to dark brown	Olivaceous to dark brown	Olivaceous to dark brown
Conidiophores	**Size (µm)**	Up to 620 µm long, 18–35 µm wide at the base, 13–18 µm wide in the middle, 7–11.5 µm wide at the apex	(430–)550–750**(–890) µm long**, 15–24(–34) µm wide at the base, 11–14(–16) µm wide in the middle, 7–10 µm(–12) wide at the apex	Up to 530 µm long, 13–15 µm wide at the base, 10–12 µm wide in the middle, 3.6–5 µm wide at the apex	160–340(–570) µm long, 12–21 µm wide at the base, 11–14 µm wide in the middle, 6–7.5 µm wide at the apex	Up to 550 µm long, 20–30 µm wide at the base, 10–17 µm wide in the middle, 5–8 µm wide at the apex	Up to 630 µm long, 18–25 µm wide at base, 10–15 µm wide in the middle, 4–7 µm wide at apex	Up to 730 µm long, 12–20 µm wide at the base, 7–14 µm wide in the middle, 5–9 µm wide at the apex
	Features	Mononematous, erect, straight or flexuous, unbranched, smooth, thick-walled, comprising 4–8 septa	Macronematous, mononematous, erect, straight or flexuous, **rarely branched**, smooth, thick-walled, wider at the base and ocasionally conical at the apex, comprising **12–17 septa**	Mononematous, basal portion sheath-like, erect, straight, unbranched, smooth, thick-walled, comprising 4–13 septa	Mononematous, erect, straight, unbranched, smooth, thick-walled, comprising 3–8 septa	Macronematous, mononematous, erect, straight or flexuous, unbranched, smooth, thick-walled, comprising 5–11 septa	Macronematous, mononematous, erect, straight or flexuous, unbranched, smooth, thick-walled, comprising 10–15 septa	Macronematous, mononematous, erect, straight or flexuous, unbranched, smooth, thick-walled, comprising 8–14 septa
	Colour	Dark brown	Brown to dark brown	Brown, becoming paler towards the apex	Brown	Not mentioned	Not mentioned	Not mentioned
Conidiogenous cells	**Size (µm)**	Not mentioned	5–12 µm long, 4–9 µm wide	7.5–13.5 × 4.2–7.4 µm	9–19 × 7.5–10.0 µm	Not mentioned	Not mentioned	Not mentioned
	Features	Holoblastic, integrated, terminal, sometimes lateral, truncate after conidium secession	Holoblastic, enteroblastic, integrated, terminal or intercalary, smooth-walled, ovoid to ampulliform, percurrently proliferating; sometimes new condiogenous cells developing on subtending cells	Flask-shaped, integrated, terminal and lateral, truncate after conidial secession, proliferating percurrently up to 4 times	Flask-shaped, integrated, terminal, truncate after conidial secession, proliferating percurrently up to 5 times	Monotretic, integrated, terminal, sometimes lateral, with up to five successive percurrent proliferations	Monotretic, integrated, terminal and lateral	Monotretic, integrated, terminal and lateral, with up to three successive percurrent proliferations
	Colour	Not mentioned	**Hyaline** or brown to dark-brown	Brown	Brown	Not mentioned	Not mentioned	Not mentioned
Conidia	**Size (µm)**	50–80 µm long, 60–90 µm wide	25–40 µm long, 30–60 µm wide	34–44 µm long, 39–52 µm wide	(73–)81–120(–135) µm long, (51–)67–94(–106) µm wide	50–65 µm long, 58–95 µm wide	27–41 µm long, 30–43 µm wide	45–50 µm long, 53–76 µm wide
	Features	Fan-shaped, lenticular in edge view, 2–3-lobed, dictyoseptate, with rows of cells radiating from a protuberant basal cell 6–7.5 µm wide	Dictyospored and cheirospored, fan-shaped to cheiroid, lenticular in edge view, occasionally apically 2–3-lobed, dictyoseptate, with up to 15 columns of cells radiating from a protuberant basal cell **2–3 µm wide**	Fan-shaped, sometimes 2–5-lobed, lenticular in edge view, dictyoseptate, with up to 15 rows of cells, basal cell protuberant, 3.7–5 µm wide	Campanulate or fan-shaped, sometimes apically 2–3-lobed, lenticular in edge view, dictyoseptate, smooth, with up to 18 vertical rows of cells, basal cell protuberant, 7–9 µm wide	Fan-shaped, sometimes 2–3-lobed, lenticular in edge view, dictyoseptate, with up to 18 rows of cells radiating from a protuberant basal cell 4.5–6.5 µm wide	Fan-shaped, sometimes lobed, lenticular in edge view, dictyoseptate, with eight rows of cells radiating from a protuberant basal cell 2.5–5 µm wide	Fan shaped, sometimes 2–3-lobed, lenticular in edge view, dictyoseptate, with 20 rows of cells radiating from a protuberant basal cell 3–5 µm wide
	Colour	Dark brown	Dark brown to olivaceous brown	Brown	Brown	Dark brown	Dark brown	Dark brown
Apical appendage(s)	**Size (µm)**	75–120 µm long, 4–6 µm wide, tapering to 1.5–2 µm wide	(2–)6–16(–20) µm long, 4–5 µm wide at the base and apex	15–30 µm long, 3–4.4 µm wide at the base, tapering to 2.4–2.8 µm wide	(33–)52–74(–126) µm long, 6.5–8.0 µm wide at the base	24–99 µm long, 3–5.5 µm wide, tapering to 1–1.5 µm wide		15–35 µm long, 3–4 µm wide, tapering to 2–3 µm wide
	Features	2–4 appendages, tapering, 3–5-septate	Without or with 1–3 appendages, rounded at the tips, **mostly untapered**, sometimes tapering 1–5-septate	2 appendages (Up to 4), tapering, 1–2-septate	2–3 appendages (rarely up to 4), tapering, 1–3(–6)-septate	2 appendages (rarely up to 3), smooth, tapering, 0–3-septate	Appendages absent	2 appendages, tapering, 1–2-septate
	Color	Brown, and apically hyaline	Pale brown to brown, becoming hyaline at the tips	Brown, and apically hyaline	Brown, and apically hyaline	Brown, and apically hyaline	-	Brown, and apically hyaline
Host(s)		On dead twigs of an unknown host	On decaying stems of *Broussonetia papyrifera*	On dried bamboo culms	On dead bamboo stem	On dead branches of *Punica granatum*	On dead branches of *Rhaphidophora decursiva*	On dead branches of *Rosa chinensis*
Distribution(s)		India	Thailand	India	India	China	China	China
References		(Bhat and Kendrick 1993)	This study	Sureshkumar et al. (2005)	Dubey and Jash (2024)	Zhang et al. (2009), Zhang et al. (2014)	Zhang et al. (2009, 2014)	Zhang et al. (2009), Zhang et al. (2014)

to *Pseudoberkleasmium*, with maximum statistical support in both methods of analysis (Fig. 1). Pleosporales taxa were accommodated in over 90 families (with and without molecular data), and have a worldwide distribution in terrestrial and aquatic habitats (Hongsanan et al. 2020; Bhunjun et al. 2021; Jiang et al. 2021; Wijayawardene et al. 2020, 2022a; Yang et al. 2022; Tang et al. 2023; Hyde et al. 2024a, b; Pem et al. 2024; Wei et al. 2024). In this study, we maintain 95 families in this order: 92 with DNA sequence data and two without (Diademaceae and Lizoniaceae). Additionally, we exclude Mycoporaceae (*Mycoporum* spp.) from the analysis due to its divergent sequences and long branches in the preliminary phylogenetic trees. Further research is needed to determine its placement within Pleosporales. Arthopyreniaceae was classified under Pleosporales in Wijayawardene et al. (2022a) and Pem et al. (2024), but we do not retain it in this order as its taxa cluster basal to Pleosporales (results not shown), similar to the findings by Thiyagaraja et al. (2021). We also retain Sublophiostomataceae within Pleosporales (Hongsanan et al. 2021), but it was not included in Pem et al. (2024). In our phylogenetic analysis, most families in Pleosporales are monophyletic, thereby further supporting their rank. A few families [e.g., Halotthiaceae (87% ML); Pseudomassarinaceae; and Testudinaceae (81% ML)] lack strong statistical support but have consistently been recovered as distinct groups in previous studies using different datasets and/or methods of analysis (Crous et al. 2018; Yang et al. 2022), suggesting that these are separate lineages distinct from other families. However, available genetic markers and DNA sequence data may not be sufficient to resolve their familial relationships with high confidence.

Xiuguozhangia is sister to *Pseudoberkleasmium* and closely related to *Hermatomyces* (Figs 1, 2). Given that we assigned *Xiuguozhangia* in Pseudoberkleasmiaceae based on the analysis of dataset 1, we analysed dataset 2 to further validate its taxonomic placement, which was based on multiple strains representing all species of *Hermatomyces* and *Pseudoberkleasmium*. Since ITS and LSU are important genetic markers to delimit fungal genera, we computed the group mean distances across these gene regions between *Hermatomyces* spp., *Pseudoberkleasmium* spp. and *Xiuguozhangia* spp. Considering the inter-generic nucleotide differences between each group (Table 2), the establishment of *Xiuguozhangia* as a distinct genus from *Pseudoberkleasmium* is supported.

The characterisation of *Xiuguozhangia* as a separate genus from *Pseudoberkleasmium* is further corroborated based on their distinct features. *Xiuguozhangia* differs from *Pseudoberkleasmium* in their colonies on natural substrates, and conidiophore, conidiogenous cell and conidial features (Zhang et al. 2014; Tibpromma et al. 2018). *Xiuguozhangia* species are depicted by dictyoseptate, campanulate, cheiroid, and brown conidia comprising multiple layers of cells, either with or without apical appendages (Zhang et al. 2014). However, *Pseudoberkleasmium* species are characterised by broadly ellipsoidal to obovoid, flattened, one-cell thick, muriform, and brown olivaceous green conidia without any apical appendage (Tibpromma et al. 2018). Conidial images of *Pseudoberkleasmium* and *Hermatomyces* are illustrated in Fig. 2. Since other species of *Xiuguozhangia* do not have molecular data, we were unable to compare the inter-species nucleotide differences, and we had to rely solely on morphology to establish our new species, *X. broussonetiae*.

With respect to the inter-species morphological differences in the genus, the new species, *X. broussonetiae*, differs from all extant species (Table 3), but exhibits overlapping features with *X. rosae*. Nonetheless, *X. broussonetiae* differs from *X. rosae* in terms of the number and characters of the apical appendages, as outlined in the note section above. Other observed differences are given in Table 3. *Xiuguozhangia* was classified as Ascomycota genus *incertae sedis* in the latest outline by Wijayawardene et al. (2022a), as it was difficult to propose its precise familial placement without the availability of DNA sequence data. Since our study is the first to provide molecular data, we refer *Xiuguozhangia* in Pseudoberkleasmiaceae. Hyde et al. (2019) initially introduced this family as monotypic to accommodate *Pseudoberkleasmium* species based on morphology and molecular data. *Pseudoberkleasmium* was described by Tibpromma et al. (2018) based on morphology and multigene phylogenetic analyses. *Pseudoberkleasmium* taxa resemble those of *Berkleasmium*, but *Berkleasmium* belongs to Tubeufiales (Tibpromma et al. 2018; Wijayawardene et al. 2022a; this study). Besides *Pseudoberkleasmium*, we treat *Xiuguozhangia* as a member of Pseudoberkleasmiaceae based on our inferred phylogenies. Upon the addition of *X. broussonetiae* to the genus, there are seven species of *Xiuguozhangia*, sharing the same holoblastic conidial ontogeny as a generic feature.

Due to the limited taxon sampling and analysis, and the lack of comprehensive DNA sequence data, there remains uncertainty in using conidial ontogeny as a basis for distinguishing *Xiuguozhangia* from *Piricaudiopsis*. As such, we refrain from drawing further conclusions regarding their taxonomic separation at this stage. Instead, a more effective approach would involve extensive sampling across diverse localities and habitats, particularly those that support the growth of hyphomycetes. Coupled with a systematic acquisition of DNA sequence data, these efforts would provide a more robust support for classifying *Xiuguozhangia* species and other poorly studied Pleosporalean taxa (Liu et al. 2024b; Pem et al. 2024). Moreover, a thorough re-examination of the type species of each genus is essential to ascertain whether these two genera are indeed distinct or if they might represent a single taxonomic entity.

Acknowledgments

We thank Dr Shaun Pennycook for suggesting and validating the nomenclature of the new species. Deecksha Gomdola thanks the Mae Fah Luang University for providing the MFU student scholarship 2020 (reference: GR-ST-FS-63-06) to pursue to PhD program, and for allocating the thesis writing grant [reference: 7702(6)/201 (No.0275)]. Deecksha Gomdola thanks the Mushroom Research Foundation (MRF) for partly funding her project, and the staffs of the MFU botanical garden for their help in the sample collection and identification of the host. Deecksha Gomdola also thanks Prof. Dr. Yong Wang of the Guizhou University (Department of Plant Pathology) for providing funds to conduct molecular work under his NFSC grant No. 31972222. We would also like to acknowledge the grant GDPX[2019]13 for partly funding this project. We thank Miss Tang Xia and Mr. Xingguo Tian for providing conidial images of *Hermatomyces* and *Pseudoberkleasmium* to complement Fig. 2. The authors extend their appreciation to the Researchers Supporting Project number (RSP2024R114), King Saud University, Riyadh, Saudi Arabia.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This work was supported by Guizhou University (NFSC grant No. 31972222 and GDPX[2019]13), Mae Fah Luang University, Mushroom Research Foundation and MFU student scholarship 2020 (reference: GR-ST-FS-63-06); thesis writing grant [reference: 7702(6)/201 (No.0275)]; Researchers Supporting Project number (RSP2024R114).

Author contributions

Conceptualization: DG, EG, RJ. Data curation: DG. Formal analysis: DG. Funding acquisition: YW. Investigation: DG. Methodology: DG. Resources: YW, FA, KDH. Software: DG. Supervision: KDH, YW, RSJ. Validation: RSJ, YW, RJ, FA, KDH, EG. Visualization: DG. Writing - original draft: DG. Writing - review and editing: RSJ, YW, RJ, KDH, EG.

Author ORCIDs

Deecksha Gomdola https://orcid.org/0000-0002-0817-1555
Rajesh Jeewon https://orcid.org/0000-0002-8563-957X
Eleni Gentekaki https://orcid.org/0000-0002-3306-6714
Ruvishika S. Jayawardena https://orcid.org/0000-0001-7702-4885
Kevin D. Hyde https://orcid.org/0000-0002-2191-0762
Fatimah Alotibi https://orcid.org/0000-0003-3629-5755

Data availability

All data generated or analysed during this study are included in this published article.

References

Bao DF, Hyde KD, McKenzie EHC, Jeewon R, Su HY, Nalumpang S, Luo ZL (2021) Biodiversity of lignicolous freshwater hyphomycetes from China and Thailand and description of sixteen species. Journal of Fungi 7(8): 669. https://doi.org/10.3390/jof7080669

Bhat DJ, Kendrick B (1993) Twenty-five new conidial fungi from the Western Ghats and the Andaman Islands (India). Mycotaxon 49: 19–90.

Bhunjun CS, Phukhamsakda C, Jeewon R, Promputtha I, Hyde KD (2021) Integrating different lines of evidence to establish a novel ascomycete genus and family (*Anastomitrabeculia*, anastomitrabeculiaceae) in pleosporales. Journal of Fungi 7(2): 1–18. https://doi.org/10.3390/jof7020094

Bhunjun CS, Niskanen T, Suwannarach N, Wannathes N, Chen YJ, McKenzie EHC, Maharachchikumbura SSN, Buyck B, Zhao CL, Fan YG, Zhang JY, Dissanayake AJ, Marasinghe DS, Jayawardena RS, Kumla J, Padamsee M, Chen YY, Liimatainen K, Ammirati JF, Phukhamsakda C, Liu JK, Phonrob W, Randrianjohany É, Hongsanan S, Cheewangkoon R, Bundhun D, Khuna S, Yu WJ, Deng LS, Lu YZ, Hyde KD, Lumyong S (2022) The numbers of fungi: Are the most speciose genera truly diverse? Fungal Diversity 114(1): 387–462. https://doi.org/10.1007/s13225-022-00501-4

Calabon MS, Hyde KD, Jones EBG, Bao DF, Bhunjun CS, Phukhamsakda C, Shen HW, Gentekaki E, Al Sharie AH, Barros J, Chandrasiri KSU, Hu DM, Hurdeal VG, Rossi W, Valle LG, Zhang H, Figueroa M, Raja HA, Seena S, Song HY, Dong W, El-Elimat T, Leonardi M, Li Y, Li YJ, Luo ZL, Ritter CD, Strongman DB, Wei MJ, Balasuriya A (2023) Freshwater fungal biology. Mycosphere 14(1): 195–413. https://doi.org/10.5943/mycosphere/14/1/4

Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T (2009) trimAl: A tool for automated alignment trimming in large-scale phylogenetic analyses. Bioinformatics 25(15): 1972–1973. https://doi.org/10.1093/bioinformatics/btp348

Carbone I, Kohn LM (1999) A method for designing primer sets for speciation studies in filamentous ascomycetes. Mycologia 91(3): 553–556. https://doi.org/10.1080/00275514.1999.12061051

Chaiwan N, Gomdola D, Wang S, Monkai J, Tibpromma S, Doilom M, Wanasinghe DN, Mortimer PE, Lumyong S, Hyde KD (2021) an online database providing updated information of microfungi in the Greater Mekong Subregion. Mycosphere 12(1): 1513–1526. https://doi.org/10.5943/mycosphere/12/1/19

Crous PW, Schumacher RK, Wingfield MJ, Akulov A, Denman S, Roux J, Braun U, Burgess TI, Carnegie AJ, Váczy KZ, Guatimosim E, Schwartsburd PB, Barreto RW, Hernández-Restrepo M, Lombard L, Groenewald JZ (2018) New and interesting Fungi. 1. Fungal Systematics and Evolution 1(1): 169–215. https://doi.org/10.3114/fuse.2018.01.08

Dong W, Hyde KD, Jeewon R, Liao CF, Zhao HJ, Kularathnage ND, Li H, Yang YH, Pem D, Shu YX, Gafforov Y, Manawasinghe IS, Doilom M (2023) Mycosphere notes 449–468: Saprobic and endophytic fungi in China, Thailand, and Uzbekistan. Mycosphere 14(1): 2208–2262. https://doi.org/10.5943/mycosphere/14/1/26

Dubey R, Jash S (2024) *Xiuguozhangia macrospora* (Pezizomycotina: Ascomycota)—a new species of bambusicolous fungi from India. Phytotaxa 644(1): 69–73. https://doi.org/10.11646/phytotaxa.644.1.10

Gilbert GS, Parker IM (2023) How to be a fungus. The Evolutionary Ecology of Plant Disease, 27_27. https://doi.org/10.1093/oso/9780198797876.003.0003

Hongsanan S, Hyde KD, Phookamsak R, Wanasinghe DN, McKenzie EHC, Sarma VV, Lücking R, Boonmee S, Bhat JD, Liu NG, Tennakoon DS, Pem D, Karunarathna A, Jiang SH, Jones GEB, Phillips AJL, Manawasinghe IS, Tibpromma S, Jayasiri SC, Sandamali D, Jayawardena RS, Wijayawardene NN, Ekanayaka AH, Jeewon R, Lu YZ, Phukhamsakda C, Dissanayake AJ, Zeng XY, Luo ZL, Tian Q, Thambugala KM, Dai D, Samarakoon MC, Chethana KWT, Ertz D, Doilom M, Liu JK, Pérez-Ortega S, Suija A, Senwanna C, Wijesinghe SN, Niranjan M, Zhang S-N, Ariyawansa HA, Jiang H-B, Zhang J-F, Norphanphoun C, de Silva NI, Thiyagaraja V, Zhang H, Bezerra JDP, Miranda-González R, Aptroot A, Kashiwadani H, Harishchandra D, Sérusiaux E, Abeywickrama PD, Bao D-F, Devadatha B, Wu H-X, Moon KH, Gueidan C, Schumm F, Bundhun D, Mapook A, Monkai J, Bhunjun CS, Chomnunti P, Suetrong S, Chaiwan N, Dayarathne MC, Yang J, Rathnayaka AR, Xu J-C, Zheng J, Liu G, Feng Y, Xie N (2020) Refined families of Dothideomycetes: Orders and families incertae sedis in Dothideomycetes. Fungal Diversity 105(1): 17–318. https://doi.org/10.1007/s13225-020-00462-6

Hongsanan S, Phookamsak R, Goonasekara ID, Thambugala KM, Hyde KD, Bhat JD, Suwannarach N, Cheewangkoon R (2021) Introducing a new pleosporalean family Sublophiostomataceae fam. nov. to accommodate *Sublophiostoma* gen. nov. Scientific Reports 11(1): 9496. https://doi.org/10.1038/s41598-021-88772-w

Hu Y, Liu J, Xu Z, Castañeda-Ruíz RF, Zhang K, Ma J (2023) Morphology and multigene phylogeny revealed three new species of *Helminthosporium* (Massarinaceae, Pleosporales) from China. Journal of Fungi 9(2): 280. https://doi.org/10.3390/jof9020280

Huelsenbeck JP, Ronquist F, Nielsen R, Bollback JP (2001) Bayesian inference of phylogeny and its impact on evolutionary biology. Science 294(5550): 2310–2314. https://doi.org/10.1126/science.1065889

Hyde KD, Tennakoon DS, Jeewon R, Bhat DJ, Maharachchikumbura SSN, Rossi W, Leonardi M, Lee HB, Mun HY, Houbraken J, Nguyen TTT, Jeon SJ, Frisvad JC, Wanasinghe DN, Lücking R, Aptroot A, Cáceres MES, Karunarathna SC, Hongsanan S, Phookamsak R, de Silva NI, Thambugala KM, Jayawardena RS, Senanayake IC, Boonmee S, Chen J, Luo ZL, Phukhamsakda C, Pereira OL, Abreu VP, Rosado AWC, Bart B, Randrianjohany E, Hofstetter V, Gibertoni TB, Soares AM da S, Plautz Jr HL, Sotão HMP, Xavier WKS, Bezerra JDP, de Oliveira TGL, de Souza-Motta CM, Magalhães OMC, Bundhun D, Harishchandra D, Manawasinghe IS, Dong W, Zhang SN, Bao DF, Samarakoon MC, Pem D, Karunarathna A, Lin CG, Yang J, Perera RH, Kumar V, Huang SK, Dayarathne MC, Ekanayaka AH, Jayasiri SC, Xiao Y, Konta S, Niskanen T, Liimatainen K, Dai YC, Ji XH, Tian XM, Mešić A, Singh SK, Phutthacharoen K, Cai L, Sorvongxay T, Thiyagaraja V, Norphanphoun C, Chaiwan N, Lu YZ, Jiang HB, Zhang JF, Abeywickrama PD, Aluthmuhandiram JVS, Brahmanage RS, Zeng M, Chethana T, Wei D, Réblová M, Fournier J, Nekvindová J, do Nascimento Barbosa R, dos Santos JEF, de Oliveira NT, Li G-J, Ertz D, Shang Q-J, Phillips AJL, Kuo C-H, Camporesi E, Bulgakov TS, Lumyong S, Jones EBG, Chomnunti P, Gentekaki E, Bungartz F, Zeng X-Y, Fryar S, Tkalčec Z, Liang J, Li G, Wen T-C, Singh PN, Gafforov Y, Promputtha I, Yasanthika E, Goonasekara ID, Zhao R-L, Zhao Q, Kirk PM, Liu J-K, Yan JY, Mortimer PE, Xu J, Doilom M (2019) Fungal diversity notes 1036–1150: Taxonomic and phylogenetic contributions on genera and species of fungal taxa. Fungal Diversity 96(1): 1–242. https://doi.org/10.1007/s13225-019-00429-2

Hyde KD, Jeewon R, Chen YJ, Bhunjun CS, Calabon MS, Jiang HB, Lin CG, Norphanphoun C, Sysouphanthong P, Pem D, Tibpromma S, Zhang Q, Doilom M, Jayawardena RS, Liu JK, Maharachchikumbura SSN, Phukhamsakda C, Phookamsak R, Al-Sadi AM, Thongklang N, Wang Y, Gafforov Y, Gareth Jones EB, Lumyong S (2020) The numbers of fungi: Is the descriptive curve flattening? Fungal Diversity 103(1): 219–271. https://doi.org/10.1007/s13225-020-00458-2

Hyde KD, Norphanphoun C, Ma J, Yang HD, Zhang JY, Du TY, Gao Y, Gomes de Farias AR, He SC, He YK, Li CJY, Li JY, Liu XF, Lu L, Su HL, Tang X, Tian XG, Wang SY, Wei DP, Xu RF, Xu RJ, Yang YY, Zhang F, Zhang Q, Bahkali AH, Boonmee S, Chethana KWT, Jayawardena RS, Lu YZ, Karunarathna SC, Tibpromma S, Wang Y, Zhao Q (2023) Mycosphere notes 387–412 – novel species of fungal taxa from around the world. Mycosphere 14(1): 663–744. https://doi.org/10.5943/mycosphere/14/1/8

Hyde KD, Wijesinghe SN, Afshari N, Aumentado HD, Bhunjun CS, Boonmee S, Camporesi E, Chethana KWT, Doilom M, Dong W, Du TY, Farias ARG, Gao Y, Jayawardena RS, Karimi O, Karunarathna SC, Kularathnage ND, Lestari AS, Li CJY, Li YX, Liao CF, Liu XF, Lu L, Lu YZ, Luo ZL, Ma J, Mamarabadi M, Manawasinghe IS, Mapook A, Mi LX, Niranjan M, Senanayake IC, Shen HW, Su HL, Tibpromma S, Xu RJ, Yan JY, Yang YH, Yang YY, Yu FQ, Kang JC, Zhang JY (2024a) Mycosphere Notes 469–520. Mycosphere 15(1): 1294–1454. https://doi.org/10.5943/mycosphere/15/1/11

Hyde KD, Wijesinghe SN, Amuhenage TB, Bera I, Bhunjun CS, Chethana KWT, De Farias ARG, Dong W, Du TY, Gao Y, Gui H, Gunarathne A, Jayawardena RS, Li H, Liu JK, Madagammana AD, Manawasinghe IS, Mapook A, Mukhopadhyay S, Niranjan M, Phukhamsakda C, Ren GC, Senanayake IC, Shu YX, Thakshila SAD, Thongklang N, Tibpromma S, Tun ZL, Wu N, Xiong YR, Xu RF, Yang YH, Yu FQ, Zhao HJ (2024b) Current Research in Applied and Environmental Mycology Fungal Profiles 1–30. Current Research in Environmental & Applied Mycology 14: 167–266. [Journal of Fungal Biology]

Hyde KD, Noorabadi MT, Thiyagaraja V, He MQ, Johnston PR, et al. (2024c) The 2024 Outline of Fungi and fungus-like taxa. Mycosphere 15(1): 5146–6239.

Index Fungorum (2024) Index Fungorum. http://www.indexfungorum.org/Names/Names.asp [accessed on 20 November 2024]

Jayasiri SC, Hyde KD, Ariyawansa HA, Bhat J, Buyck B, Cai L, Dai YC, Abd-Elsalam KA, Ertz D, Hidayat I, Jeewon R, Jones EBG, Bahkali AH, Karunarathna SC, Liu JK, Luangsa-ard JJ, Lumbsch HT, Maharachchikumbura SSN, McKenzie EHC, Moncalvo JM, Ghobad-Nejhad M, Nilsson H, Pang KL, Pereira OL, Phillips AJL, Raspé O, Rollins AW, Romero AI, Etayo J, Selçuk F, Stephenson SL, Suetrong S, Taylor JE, Tsui CKM, Vizzini A, Abdel-Wahab MA, Wen TC, Boonmee S, Dai DQ, Daranagama DA, Dissanayake AJ, Ekanayaka AH, Fryar SC, Hongsanan S, Jayawardena RS, Li WJ, Perera RH, Phookamsak R, de Silva NI, Thambugala KM, Tian Q, Wijayawardene NN, Zhao RL, Zhao Q, Kang JC, Promputtha I (2015) The Faces of Fungi database: Fungal names linked with morphology, phylogeny and human impacts. Fungal Diversity 74(1): 3–18. https://doi.org/10.1007/s13225-015-0351-8

Jayawardena RS, Hyde KD, Wang S, Sun YR, Suwannarach N, Sysouphanthong P, Abdel-Wahab MA, Abdel-Aziz FA, Abeywickrama PD, Abreu VP, Armand A, Aptroot A, Bao DF, Begerow D, Bellanger JM, Bezerra JDP, Bundhun D, Calabon MS, Cao T, Cantillo T, Carvalho JLVR, Chaiwan N, Chen CC, Courtecuisse R, Cui BK, Damm U, Denchev CM, Denchev TT, Deng CY, Devadatha B, de Silva NI, dos Santos LA, Dubey NK, Dumez S, Ferdinandez HS, Firmino AL, Gafforov Y, Gajanayake AJ, Gomdola D, Gunaseelan S, Shucheng-He, Htet ZH, Kaliyaperumal M, Kemler M, Kezo K, Kularathnage ND, Leonardi M, Li J-P, Liao C, Liu S, Loizides M, Luangharn T, Ma J, Madrid H, Mahadevakumar S, Maharachchikumbura SSN, Manamgoda DS, Martín MP, Mekala N, Moreau P-A, Mu Y-H, Pahoua P, Pem D, Pereira OL, Phonrob W, Phukhamsakda C, Raza M, Ren G-C, Rinaldi AC, Rossi W, Samarakoon BC, Samarakoon MC, Sarma VV, Senanayake IC, Singh A, Souza MF, Souza-Motta CM, Spielmann AA, Su W, Tang X, Tian XG, Thambugala KM, Thongklang N, Tennakoon DS, Wannathes N, Wei DP, Welti S, Wijesinghe SN, Yang H, Yang Y, Yuan H-S, Zhang H, Zhang J, Balasuriya A, Bhunjun CS, Bulgakov TS, Cai L, Camporesi E, Chomnunti P, Deepika YS, Doilom M, Duan W-J, Han S-L, Huanraluek N, Jones EBG, Lakshmidevi N, Li Y, Lumyong S, Luo Z-L, Khuna S, Kumla J, Manawasinghe IS, Mapook A, Punyaboon W, Tibpromma S, Lu Y-Z, Yan JY, Wang Y (2023) Fungal diversity notes 1512–1610: Taxonomic and phylogenetic contributions on genera and species of fungal taxa. Fungal Diversity 117(1): 1–272. https://doi.org/10.1007/s13225-022-00513-0

Jiang HB, Jeewon R, Karunarathna SC, Phukhamsakda C, Doilom M, Kakumyan P, Suwannarach N, Phookamsak R, Lumyong S (2021) Reappraisal of Immotthia in Dictyosporiaceae, Pleosporales: Introducing *Immotthia bambusae* sp. nov. and *Pseudocyclothyriella clematidis* comb. et gen. nov. based on morphology and phylogeny. Frontiers in Microbiology 12: 656235. https://doi.org/10.3389/fmicb.2021.656235

Katoh K, Rozewicki J, Yamada KD (2019) MAFFT online service: Multiple sequence alignment, interactive sequence choice and visualization. Briefings in Bioinformatics 20(4): 1160–1166. https://doi.org/10.1093/bib/bbx108

Liu Y, Whelen S, Hall BD (1999) Phylogenetic relationships among ascomycetes: Evidence from an RNA polymerse II subunit. Molecular Biology and Evolution 16(12): 1799–1808. https://doi.org/10.1093/oxfordjournals.molbev.a026092

Liu SL, Wang XW, Li GJ, Deng CY, Rossi W, Leonardi M, Liimatainen K, Kekki T, Niskanen T, Smith ME, Ammirati J, Bojantchev D, Abdel-Wahab MA, Zhang M, Tian E, Lu YZ, Zhang JY, Ma J, Dutta AK, Acharya K, Du TY, Xu J, Kim JS, Lim YW, Gerlach A, Zeng NK, Han

YX, Razaghi P, Raza M, Cai L, Calabon MS, Jones EBG, Saha R, Kumar TKA, Krishnapriya K, Thomas A, Kaliyaperumal M, Kezo K, Gunaseelan S, Singh SK, Singh PN, Lagashetti AC, Pawar KS, Jiang S, Zhang C, Zhang H, Qing Y, Bau T, Peng XC, Wen TC, Ramirez NA, Niveiro N, Li MX, Yang ZL, Wu G, Tarafder E, Tennakoon DS, Kuo CH, da Silva TM, Souza-Motta CM, Bezerra JDP, He G, Ji XH, Suwannarach N, Kumla J, Lumyong S, Wannathes N, Rana S, Hyde KD, Zhou LW (2024a) Fungal diversity notes 1717–1817: Taxonomic and phylogenetic contributions on genera and species of fungal taxa. Fungal Diversity 124(1): 1–216. https://doi.org/10.1007/s13225-023-00529-0

Liu NG, Hyde KD, Sun YR, Bhat DJ, Jones EBG, Jumpathong J, Lin CG, Lu YZ, Yang J, Liu LL, Liu ZY, Liu JK (2024b) Notes, outline, taxonomy and phylogeny of brown-spored hyphomycetes. Fungal Diversity 128: 1–282. https://doi.org/10.1007/s13225-023-00529-0

Ma J, Gomdola D, Boonmee S, Shen HW, Tang X, Zhang LJ, Lu YZ, Hyde KD (2024) Three new species of *Neohelicomyces* (Tubeufiales, Tubeufiaceae) from freshwater and terrestrial habitats in China. MycoKeys 105: 317–336. https://doi.org/10.3897/mycokeys.105.124129

Maharachchikumbura SSN, Chen Y, Ariyawansa HA, Hyde KD, Haelewaters D, Perera RH, Samarakoon MC, Wanasinghe DN, Bustamante DE, Liu JK, Lawrence DP, Cheewangkoon R, Stadler M (2021) Integrative approaches for species delimitation in Ascomycota. Fungal Diversity 109(1): 155–179. https://doi.org/10.1007/s13225-021-00486-6

Manawasinghe IS, Calabon MS, Jones EBG, Zhang YX, Liao CF, Xiong Y, Chaiwan N, Kularathnage ND, Liu NG, Tang SM, Sysouphanthong P, Du TY, Luo M, Pasouvang P, Pem D, Phonemany M, Ishaq M, Chen JW, Karunarathna SC, Mai ZL, Rathnayaka AR, Samarakoon MC, Tennakoon DS, Wijesinghe SN, Yang YH, Zhao HJ, Fiaz M, Doilom M, Dutta AK, Khalid AN, Liu JW, Thongklang N, Senanayake IC, Tibpromma S, You LQ, Camporesi E, Gafforov YS, Hyde KD (2022) Mycosphere notes 345–386. Mycosphere 13(1): 454–557. https://doi.org/10.5943/mycosphere/13/1/3

Mena PJ, Mercado SA (1987) *Piricaudiopsis* (Hyphomycetes, Deuteromycotina), nuevo género enteroblástico de Cuba. Acta Botanica Cubana 51: 1–5.

Miller MA, Pfeiffer W, Schwartz T (2010) Creating the CIPRES Science Gateway for inference of large phylogenetic trees. Proceedings of the 2011 TeraGrid Conference: extreme digital discovery, 8 pp. https://doi.org/10.1145/2016741.2016785

Nguyen LT, Schmidt HA, Von Haeseler A, Minh BQ (2015) IQ-TREE: A fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. Molecular Biology and Evolution 32(1): 268–274. https://doi.org/10.1093/molbev/msu300

O'Donnell K, Kistlerr HC, Cigelnik E, Ploetz RC (1998) Multiple evolutionary origins of the fungus causing panama disease of banana: Concordant evidence from nuclear and mitochondrial gene genealogies. Proceedings of the National Academy of Sciences of the United States of America 95(5): 2044–2049. https://doi.org/10.1073/pnas.95.5.2044

Pem D, Jeewon R, Chethana KWT, Hongsanan S, Doilom M, Suwannarach N, Hyde KD (2021) Species concepts of Dothideomycetes: Classification, phylogenetic inconsistencies and taxonomic standardization. Fungal Diversity 109(1): 283–319. https://doi.org/10.1007/s13225-021-00485-7

Pem D, Hyde KD, McKenzie EHC, Hongsanan S, Wanasinghe DN, Boonmee S, Darmostuk V, Bhat JD, Tian Q, Htet ZH, Senanayake IC, Niranjan M, Sarma VV, Doilom M, Dong W (2024) A comprehensive overview of genera in Dothideomycetes. Mycosphere 15(1): 2175–4568. https://doi.org/10.5943/mycosphere/15/1/18

Perera RH, Hyde KD, Jones EBG, Maharachchikumbura SSN, Bundhun D, Camporesi E, Akulov A, Liu JK, Liu ZY (2023) Profile of Bionectriaceae, Calcarisporiaceae, Hypocre-

aceae, Nectriaceae, Tilachlidiaceae, Ijuhyaceae fam. nov., Stromatonectriaceae fam. nov. and Xanthonectriaceae fam. nov. Fungal Diversity 118(1): 95–271. https://doi.org/10.1007/s13225-022-00512-1

Rambaut A, Drummond A (2014) FigTree v1. 3.1 Institute of Evolutionary Biology. University of Edinburgh.

Rathnayaka AR, Tennakoon DS, Jones GE, Wanasinghe DN, Bhat DJ, Priyashantha AH, Stephenson SL, Tibpromma S, Karunarathna SC (2024) Significance of precise documentation of hosts and geospatial data of fungal collections, with an emphasis on plant-associated fungi. New Zealand Journal of Botany 31: 1–28. https://doi.org/10.1080/0028825X.2024.2381734

Ronquist F, Huelsenbeck JP (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. Bioinformatics 19(12): 1572–1574. https://doi.org/10.1093/bioinformatics/btg180

Seifert K, Morgan-Jones G, Gams W, Kendrick B (2011) The genera of hyphomycetes. CBS Biodiversity Series 9: 1–997.

Senanayake IC, Rathnayaka AR, Marasinghe DS, Calabon MS, Gentekaki E, Lee HB, Hurdeal VG, Pem D, Dissanayake LS, Wijesinghe SN, Bundhun D, Nguyen TT, Goonasekara ID, Abeywickrama PD, Bhunjun CS, Jayawardena RS, Wanasinghe DN, Jeewon R, Bhat DJ, Xiang MM (2020) Morphological approaches in studying fungi: Collection, examination, isolation, sporulation and preservation. Mycosphere 11(1): 2678–2754. https://doi.org/10.5943/mycosphere/11/1/20

Senanayake IC, Rossi W, Leonardi M, Weir A, McHugh M, Rajeshkumar KC, Verma RK, Karunarathna SC, Tibpromma S, Ashtekar N, Ashtamoorthy SK, Raveendran S, Kour G, Singh A, De la Peña-Lastra S, Mateos A, Kolařík M, Antonín V, Ševčíková H, Esteve-Raventós F, Larsson E, Pancorbo F, Moreno G, Altés A, Turégano Y, Du TY, Lu L, Li QR, Kang JC, Gunaseelan S, Kezo K, Kaliyaperumal M, Fu J, Samarakoon MC, Gafforov Y, Teshaboeva S, Kunjan PC, Chamaparambath A, Flakus A, Etayo J, Rodriguez-Flakus P, Zhurbenko MP, de Silva NI, Tennakoon DS, Latha KPD, Manimohan P, Raj KNA, Calabon MS, Ahmadpour A, Heidarian Z, Alavi Z, Alavi F, Ghosta Y, Azizi R, Luo M, Zhao MP, Kularathnage ND, Hua L, Yang YH, Liao CF, Zhao HJ, Lestari AS, Jayasiri SC, Yu FM, Lei L, Liu JW, Karimi O, Tang SM, Sun YR, Wang Y, Zeng M, Htet ZH, Linaldeddu BT, Alves A, Phillips AJL, Bregant C, Montecchio L, De Kesel A, Hustad VP, Miller AN, Fedosova AG, Kučera V, Raza M, Hussain M, Chen YP, Thiyagaraja V, Gomdola D, Rathnayaka AR, Dissanayake AJ, Suwannarach N, Hongsanan S, Maharachchikumbura SSN, Dissanayake LS, Wijayawardene NN, Phookamsak R, Lumyong S, Jones EBG, Yapa N, Wanasinghe DN, Xie N, Doilom M, Manawasinghe IS, Liu JK, Zhao Q, Xu B, Hyde KD, Song J (2023) Fungal diversity notes 1611–1716: Taxonomic and phylogenetic contributions on fungal genera and species emphasis in south China. Fungal Diversity 122(1): 161–403. https://doi.org/10.1007/s13225-023-00523-6

Shenoy BD, Jeewon R, Wu WP, Bhat DJ, Hyde KD (2006) Ribosomal and RPB2 DNA sequence analyses suggest that *Sporidesmium* and morphologically similar genera are polyphyletic. Mycological Research 110(8): 916–928. https://doi.org/10.1016/j.mycres.2006.06.004

Sureshkumar G, Sharath Babu K, Kunwar IK, Manoharachary C (2005) Two new hyphomycetous fungal species from India. Mycotaxon 92: 279–283.

Tamura K, Peterson D, Stecher G, Filipski A, Kumar S (2013) MEGA6: Molecular Evolutionary Genetics Analysis Version 6.0. Molecular Biology and Evolution 30(12): 2725–2729. https://doi.org/10.1093/molbev/mst197

Tang X, Jeewon R, Lu YZ, Alrefaei AF, Jayawardena RS, Xu RJ, Ma J, Chen XM, Kang JC (2023) Morphophylogenetic evidence reveals four new fungal species within Tetra-

plosphaeriaceae (Pleosporales, Ascomycota) from tropical and subtropical forest in China. MycoKeys 100: 153–170. https://doi.org/10.3897/mycokeys.100.113141

Thiyagaraja V, Lücking R, Ertz D, Coppins BJ, Wanasinghe DN, Karunarathna SC, Suwannarach N, To-Anun C, Cheewangkoon R, Hyde KD (2021) Sequencing of the type species of *Arthopyrenia* places Arthopyreniaceae as a synonym of Trypetheliaceae. Mycosphere 12(1): 993–1011. https://doi.org/10.5943/mycosphere/12/1/10

Tian XG, Bao DF, Karunarathna SC, Jayawardena RS, Hyde KD, Bhat DJ, Luo ZL, Elgorban AM, Hongsanan S, Rajeshkumar KC, Maharachchikumbura SSN, Suwannarach N, Dawoud TM, Lu YZ, Han JJ, Xiao YP, Du TY, Lu L, Xu RF, Dai DQ, Liu XF, Liu C, Tibpromma S (2024) Taxonomy and phylogeny of Ascomycetes associated with selected economically important monocotyledons in China and Thailand. Mycosphere 15(1): 1–274. https://doi.org/10.5943/mycosphere/15/1/1

Tibpromma S, Hyde KD, McKenzie EHC, Bhat DJ, Phillips AJL, Wanasinghe DN, Samarakoon MC, Jayawardena RS, Dissanayake AJ, Tennakoon DS, Doilom M, Phookamsak R, Tang AMC, Xu J, Mortimer PE, Promputtha I, Maharachchikumbura SSN, Khan S, Karunarathna SC (2018) Fungal diversity notes 840–928: Micro-fungi associated with Pandanaceae. Fungal Diversity 93(1): 1–160. https://doi.org/10.1007/s13225-018-0408-6

Vaidya G, Lohman DJ, Meier R (2011) SequenceMatrix: Concatenation software for the fast assembly of multi-gene datasets with character set and codon information. Cladistics 27(2): 171–180. https://doi.org/10.1111/j.1096-0031.2010.00329.x

Wei GY, Mo WD, Hu Y, Zou MT, Wen JT, Chang LF, Chen YS, Norphanphoun C, Yang QI, Wang Y (2024) – Discovery of a Novel *Occultibambusa* Species (Occultibambusaceae, Dothideomycetes) from Poaceae Host from Guizhou Province, China. Current Research in Environmental & Applied Mycology 14: 146–156. [Journal of Fungal Biology]

White TJ, Bruns T, Lee SJWT, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. PCR Protocols 64: 315–322. https://doi.org/10.1016/B978-0-12-372180-8.50042-1

Wijayawardene NN, Hyde KD, Al-Ani LKT, Tedersoo L, Haelewaters D, Rajeshkumar KC, Zhao RL, Aptroot A, Leontyev DV, Saxena RK, Tokarev YS, Dai DQ, Letcher PM, Stephenson SL, Ertz D, Lumbsch HT, Kukwa M, Issi IV, Madrid H, Phillips AJL, Selbmann L, Pfliegler WP, Horváth E, Bensch K, Kirk PM, Kolaříková K, Raja HA, Radek R, Papp V, Dima V, Ma J, Malosso E, Takamatsu S, Rambold G, Gannibal PB, Triebel D, Gautam AK, Avasthi S, Suetrong S, Timdal E, Fryar SC, Delgado G, Réblová M, Doilom M, Dolatabadi S, Pawłowska J, Humber RA, Kodsueb R, Sánchez-Castro I, Goto BT, Silva DKA, de Souza FA, Oehl F, da Silva GA, Silva IR, Błaszkowski J, Jobim K, Maia LC, Barbosa FR, Fiuza PO, Divakar PK, Shenoy BD, Castañeda-Ruiz RF, Somrithipol S, Lateef AA, Karunarathna SC, Tibpromma S, Mortimer PE, Wanasinghe DN, Phookamsak R, Xu J, Wang Y, Tian F, Alvarado P, Li DW, Kušan I, Matočec N, Maharachchikumbura SSN, Papizadeh M, Heredia G, Wartchow F, Bakhshi M, Boehm E, Youssef N, Hustad VP, Lawrey JD, Santiago ALCMA, Bezerra JDP, Souza-Motta CM, Firmino AL, Tian Q, Houbraken J, Hongsanan S, Tanaka K, Dissanayake AJ, Monteiro JS, Grossart HP, Suija A, Weerakoon G, Etayo J, Tsurykau A, Vázquez V, Mungai P, Damm U, Li QR, Zhang H, Boonmee S, Lu YZ, Becerra AG, Kendrick B, Brearley FQ, Motiejūnaitė J, Sharma B, Khare R, Gaikwad S, Wijesundara DSA, Tang LZ, He MQ, Flakus A, Rodriguez-Flakus P, Zhurbenko MP, McKenzie EHC, Stadler M, Bhat DJ, Liu JK, Raza M, Jeewon R, Nassonova ES, Prieto M, Jayalal RGU, Erdoğdu M, Yurkov A, Schnittler M, Shchepin ON, Novozhilov YK, Silva-Filho AGS, Liu P, Cavender JC, Kang Y, Mohammad S, Zhang LF, Xu RF, Li YM, Dayarathne MC, Ekanayaka AH, Wen TC, Deng CY, Pereira OL, Navathe S, Hawksworth DL, Fan XL, Dissanayake LS, Kuhnert E, Grossart HP, Thines M (2020)

Outline of Fungi and fungus-like taxa. Mycosphere 11(1): 1060–1456. https://doi.org/10.5943/mycosphere/11/1/8

Wijayawardene NN, Hyde KD, Dai DQ, Sánchez-García M, Goto BT, Saxena RK, Erdoğdu M, Selçuk F, Rajeshkumar KC, Aptroot A, Błaszkowski J, Boonyuen N, da Silva GA, de Souza FA, Dong W, Ertz D, Haelewaters D, Jones EBG, Karunarathna SC, Kirk PM, Kukwa M, Kumla J, Leontyev DV, Lumbsch HT, Maharachchikumbura SSN, Marguno F, Martínez-Rodríguez P, Mešić A, Monteiro JS, Oehl F, Pawłowska J, Pem D, Pfliegler WP, Phillips AJL, Pošta A, He MQ, Li JX, Raza M, Sruthi OP, Suetrong S, Suwannarach N, Tedersoo L, Thiyagaraja V, Tibpromma S, Tkalčec Z, Tokarev YS, Wanasinghe DN, Wijesundara DSA, Wimalaseana SDMK, Madrid H, Zhang GQ, Gao Y, Sánchez-Castro I, Tang LZ, Stadler M, Yurkov A, Thines M (2022a) – Outline of Fungi and fungus-like taxa – 2021. Mycosphere 13(1): 53–453. https://doi.org/10.5943/mycosphere/13/1/2

Wijayawardene NN, Phillips AJL, Pereira DS, Dai DQ, Aptroot A, Monteiro JS, Druzhinina IS, Cai F, Fan X, Selbmann L, Coleine C, Castañeda-Ruiz RF, Kukwa M, Flakus A, Fiuza PO, Kirk PM, Kumar KCR, Ieperuma Arachchi IS, Suwannarach N, Tang L-Z, Boekhout T, Tan CS, Jayasinghe RPPK, Thines M (2022b) Forecasting the number of species of asexually reproducing fungi (Ascomycota and Basidiomycota). Fungal Diversity 114(1): 463–490. https://doi.org/10.1007/s13225-022-00500-5

Wu W, Diao Y (2022) Anamorphic chaetosphaeriaceous fungi from China. Fungal Diversity 116(1): 1–546. https://doi.org/10.1007/s13225-022-00509-w

Wu W, Diao Y (2023) The chalara-like anamorphs of Leotiomycetes. Fungal Diversity 119(1): 213–490. https://doi.org/10.1007/s13225-023-00515-6

Yang EF, Tibpromma S, Karunarathna SC, Phookamsak R, Xu JC, Zhao ZX, Karunanayake C, Promputtha I (2022) Taxonomy and phylogeny of novel and extant taxa in Pleosporales associated with *Mangifera indica* from Yunnan, China (Series I). Journal of Fungi 8(2): 152. https://doi.org/10.3390/jof8020152

Yang J, Liu LL, Jones EBG, Hyde KD, Liu ZY, Bao DF, Liu NG, Li WL, Shen HW, Yu XD, Liu JK (2023) Freshwater fungi from karst landscapes in China and Thailand. Fungal Diversity 119(1): 1–212. https://doi.org/10.1007/s13225-023-00514-7

Zhang K, Ma J, Wang Y, Zhang XG (2009) Three new species of *Piricaudiopsis* from southern China. Mycologia 101(3): 417–422. https://doi.org/10.3852/08-147

Zhang K, Ma L, Ma J, Castañeda-Ruiz RF (2014) *Xiuguozhangia*, a new genus of microfungi to accommodate five *Piricaudiopsis* species. Mycotaxon 128(1): 131–135. https://doi.org/10.5248/128.131

Zhang SN, Hyde KD, Jones EBG, Yu XD, Cheewangkoon R, Liu JK (2024) Current insights into palm fungi with emphasis on taxonomy and phylogeny. Fungal Diversity 127(1): 55–301. https://doi.org/10.1007/s13225-024-00536-9